



Antigen Retrieval Solution for IHC, Tris EDTA- pH 9.0

Storage: 2-8°C

Catalog No: B06C-1000 1000ml

Kit Format:

1000mL liquid form.

Catalog Number	Description	Size	pH
B06C-1000	10 mM Tris 1 mM EDTA, pH 9.0 (RTU)	1000mL	9.0

Recommended Protocol:

10 mM Tris 1 mM EDTA, pH 9.0, ready-to-use buffer. Refer to HIER Procedure for details.

HIER Procedure for FFPE Tissue Sections:

1. Remove paraffin in xylene and rehydrate tissue in graded alcohols.
2. Rinse for 5 min in distilled water.
3. Place slides in a beaker containing a ready-to-use buffer. Heat it in a pressure cooker at 120°C for 2.5 min.
4. Remove the beaker with slides from the cooker and allow it cool for 25-30 minutes.
5. Block endogenous peroxidase with 3% hydrogen peroxide for 15 minutes.
6. Wash slides twice with 1X PBST (2 min each).
7. Start the immunostaining procedure.

Applications tested

Antigen Retrieval Solution for IHC, Tris EDTA - pH 9.0 has been tested by immunohistochemistry using FFPE tissue sections.

Precaution:

Wear gloves and take other necessary laboratory safety procedures.

Storage:

Store at 2-8°C. Do not freeze.

Remarks:

For research use only.