

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for 73-086

Kcnmb4 Mouse Monoclonal Antibody [Clone ID: L18A/3]

Product data:

Product Type: Primary Antibodies

Clone Name: L18A/3

Applications: IHC, IP, WB

Recommend Dilution: Immunoblot (IB)

Immunohistochemistry (IHC) Immunoprecipitation (IP)

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Fusion protein amino acids 45-167 (entire extracellular domain) of mouse BKBeta4

(accession number NP_067427).

Rat: 100% identity (123/123 amino acids identical). Human: 95% identity (117/123 amino acids identical).

Specificity: No cross-reactivity against BKBeta1, BKBeta2 or BKBeta3

Formulation: State: Supernatant

Gene Name: Mus musculus potassium large conductance calcium-activated channel, subfamily M, beta

member 4 (Kcnmb4)

Database Link: Entrez Gene 27345 HumanEntrez Gene 66016 RatEntrez Gene 58802 Mouse

Synonyms: Maxi K channel subunit beta-4, BK channel subunit beta-4, K(VCA)beta-4, Slo-beta-4,

Charybdotoxin receptor subunit beta-4






Note:

USERS will cite the UC Davis/NIH NeuroMab Facility in any publication(s) describing the research utilizing the MATERIALS. The suggested acknowledgment statement is as follows: "The monoclonal antibody _ was developed by and/or obtained from the UC Davis/NIH NeuroMab Facility, supported by NIH grant U24NS050606 and maintained by the Department of Neurobiology, Physiology and Behavior, College of Biological Sciences, University of California, Davis, CA 95616."


Also, please include the complete clone number (e.g., N52A/42) and the Antibody Registry identification number (e.g., RRID:AB_2120479) to avoid ambiguity.

View Research License Agreement

Product images:



Adult rat hippocampus immunohistochemistry



Transfected cell immunoblot: extracts of COS-1 cells transiently transfected with BKBeta4 or Kv2.1 plasmids and probed with L18A/3 pure IgG at 10 μ g/mL (left panel) or K89/41 TC supe (right panel).