

Polink-2 AP Plus Mouse-NR (No cross react to rat) Detection System for Immunohistochemistry

(2-step Polymer-AP detection system for mouse primary antibody, biotin-free,)
Polymer Detection System with Super Sensitivity and Specificity
Super clean when detecting mouse antibody on Human & Rat tissue

Storage: 4-8°C

| | | | |
|-------------|--------------------------|---------|-----------------------------|
| Catalog No. | <input type="checkbox"/> | D65-110 | 110mL (Bulk, w/o chromogen) |
| | <input type="checkbox"/> | D65-18 | 18mL |
| | <input type="checkbox"/> | D65-6 | 6mL |

Intended Use:

Detecting MOUSE primary antibody on RAT tissue is a very difficult task in research field due to background staining issues. Polink-2 Plus AP Mouse-NR (no cross react to rat) Detection Kit is specially designed to solve the problem. The secondary antibody is adsorbed to rat, rabbit and human serum proteins. This technology provides excellent specificity to detect mouse primary antibody (user supplied) on rat tissue. Besides rat tissue Polink-2 Plus AP Mouse-NR Detection kit also can be used on human tissue and rabbit tissue as well.

Polink-2 Plus AP Mouse-NR Detection Kit is the 3rd generation of polymer detection system. It uses mouse antibody enhancer to help amplify the polymer-enzyme conjugate reaction to achieve super sensitivity and specificity in immunohistochemistry staining. It produces consistent immunostaining outcomes on archival tissues and on difficult-to-work antibodies. User may need to further dilute primary antibody due to super sensitivity of Polink-2 Plus detection system. It is a biotin-free system; therefore it overcomes the non-specific staining caused by Streptavidin/biotin system due to endogenous biotin. Most commonly used specimens for this system are: frozen tissue, paraffin-embedded tissue, freshly prepared lymphocytes and fixed culture cells. It can be used for manual stain or autostainer. Staining conditions need to be optimized by user.

Polink-2 Plus AP Detection System offers a wide choice for primary antibodies, including broad spectrum (for mouse and rabbit primary antibodies), mouse, rabbit, goat, and rat primary antibodies. Refer to **Related Product** section for details.

Kit components:

| Component No. | Content | 6mL Kit | 18mL Kit | 110mL Kit |
|-------------------|------------------------------------|---------|----------|-----------|
| Reagent 1 | Mouse-NR Antibody Enhancer(RTU) | 6mL | 18mL | 110mL |
| Reagent 2 | Polymer AP for Mouse-NR (RTU) | 6mL | 18mL | 110mL |
| Reagent 3A | GBI-Permanent Red Substrate (RTU) | 7mL | 18mL | NA |
| Reagent 3B | GBI-Permanent Red Activator (5x) | 1.4mL | 1.8mLx2 | NA |
| Reagent 3C | GBI-Permanent Red Chromogen (100x) | 70µL | 180µL | NA |

Recommended Protocol:

1. Fixation: To ensure the quality of the staining and obtain reproducible performance, user needs to supply appropriately fixed tissue and well prepared slides.
2. Tissue need to be adhered to the slide tightly to avoid tissue falling off.
3. Paraffin embedded section must be deparaffinized with xylene and rehydrated with a graded series of ethanol before staining.
4. Cell smear samples should be made into as thin monolayer as possible to obtain satisfactory results.
5. Investigator needs to optimize dilution and incubation times for primary antibodies.
6. Three control slides will aid the interpretation of the result: positive tissue control, reagent control (slides treated with Isotype control reagent), and negative control.
7. Staining steps: DO NOT let specimen or tissue dry from this point on.
8. We recommend TBS-T to be used as the wash buffer to get the highest sensitivity and clean background. Phosphate in the PBS-T may inhibit the activity of the alkaline phosphatase. **Note: 1X TBS-T =50mM Tris HCl, 150mM NaCl, 0.05% Tween-20 pH7.6.** GBI sells 10xTBS-T for your convenience (B11xx)

| Reagent | Staining Procedure | Incubation Time |
|---|---|-----------------|
| 1. HIER PRETREATMENT: | a. Heat Induced Epitope Retrieval (HIER) may be required for primary antibody. Refer to primary antibody datasheet. b. Wash with PBS-T containing 0.05% Tween-20 or 1X TBS-T(See note 8 above) ; 3 times for 2 minutes each. | |
| 2. PRIMARY ANTIBODY Supplied by user | a. Apply 2 drops (100µL) or enough volume of PRIMARY ANTIBODY to cover the tissue section completely. Incubate in moist chamber for 30-60 min. b. Wash with PBS-T containing 0.05% Tween-20 or 1X TBS-T ; 3 times for 2 minutes each. | 30-60min |
| 3. Reagent 1 | a. Apply 2 drops (100µL) or enough volume of Reagent 1 Mouse-NR Antibody Enhancer to | 10min |

| | | |
|---|---|-------------------------|
| Mouse-NR Antibody Enhancer (RTU) | cover each section. Incubate in moist chamber for 10 min. b. Wash with 1X TBS-T only ; 3 times for 2 minutes each. | |
| 4. Reagent 2 Polymer AP for Mouse-NR (RTU) | a. Apply 2 drops (100µL) or enough volume of POLYMER-AP for Mouse-NR to cover each section. Incubate in moist chamber for 10 min. b. Wash with 1xTBS-T 3 times for 2 minutes each. | 10min |
| 5. Reagent 3A, 3B, 3C Reagent 3A: GBI-Permanent Red Substrate (RTU) Reagent 3B: GBI-Permanent Red Activator (5x) Reagent 3C: GBI-Permanent Red Chromogen (100x) | a. Add 200µL of Reagent 3B (Activator) into 1mL of Reagent 3A (Substrate) and mix well. Add 10µL of Reagent 3C (Chromogen) into the mixture and mix well. [Note: For fewer slides , Add 100µL of Reagent 3B (Activator) into 500µL of Reagent 3A (Substrate) and mix well. Add 5µL of Reagent 3C (Chromogen) into the mixture and mix well.] b. Apply 2 drops (100µL) or enough volume of GBI-Permanent Red working solution to completely cover the tissue. Incubate for 10 min, observe appropriate color development. c. Rinse well with distilled water. | 10min |
| 6. Hematoxylin: Supplied by user. | a. Counterstain with 2 (100uL) or more drops hematoxylin to cover tissue completely and wait about 20 seconds . b. Rinse well with tap water for 1-2 min. c. Put slides in PBS until the color turn blue (about ½ - 1 min.) d. Rinse in distill water, then rinse well with tap water | 20-30 seconds |
| 7. Mounting medium: Supplied by user | Follow the manufacture data sheet procedure for mounting. Recommended product: 1. GB-Mount: Cat. No. E01-18 (18mL), for alcohol soluble substrates (AEC, GBI-Permanent Red and AP-Blue) 2. Simpo-Mount: Cat.No. E03-18 (18mL), E03-100 (100mL), universal permanent mounting medium. Can be used with or without cover slip | Refer to insert |

Protocol Notes:

- The fixation, tissue slide thickness, and primary antibody dilution and incubation time effect results significantly. Investigator needs to consider all factors and determine optimal conditions when interpret the result.
- Pre-antibody blocking is optional and can be omitted if primary antibodies are diluted in buffers containing 2-10% normal goat serum.
- GBI-Permanent Red** is insoluble in organic solvent and can be coverslipped as well. however the dehydration steps must be shorter for optimal tissue structure and chromogen signal maintenance.

Note: Please wipe off extra water and air dry slides before dehydration and clear.

- 1x 80% Ethanol 20 seconds;
- 1x 95% Ethanol 20 seconds;
- 3x 100% Ethanol 20 seconds each;
- 1x 100% Xylene 20 seconds;
- Add 1 drop of xylene based mountant (Cat. No. O-Mount, E02-18) and coverslip. Press to push the air bubble out.

CAUTION: DO NOT dehydrate in xylene longer than 20 seconds! It will erase GBI-Permanent Red stain!

Related Products:

| Product | Catalog No. | Size | Product | Catalog No. | Size |
|---|----------------|------------|-------------------------------------|-----------------|--------------|
| Polink-2 Plus AP Broad Bulk Kit | D68-110 | 110mL | Polink-2 Plus AP Mouse Bulk Kit | D69-110 | 110mL |
| Polink-2 Plus AP Broad 18mL Kit / 6mL Kit | D68-18 / D68-6 | 18mL / 6mL | Polink-2 Plus AP Mouse 18mL/6mL Kit | D69-18 / D69-6 | 18mL / 6mL |
| Polink-2 Plus AP Rabbit Bulk Kit | D70-110 | 110mL | Fast Red Kit | C03-60 | 60mL |
| Polink-2 Plus AP Rabbit 18mL Kit / 6mL Kit | D70-18 / D70-6 | 18mL / 6mL | GBI-Permanent Red Kit | C13-18/C13-120 | 18mL/120mL |
| Polink-2 Plus AP Goat bulk | D66-110 | 110mL | BCIP/NBT Kit | C05-100/C05-18 | 100mL / 18mL |
| Polink-2 Plus AP Goat 18mL Kit / 6mL Kit | D66-18 / D66-6 | 18mL / 6mL | GB-Mount (Aqueous) | E01-18 | 18mL |
| Polink-2 Plus AP Rat-NM bulk Kit (no cross react to mouse) | D67-110 | 110mL | Simpo-Mount (Aqueous) | E03-100 /E03-18 | 100mL / 18mL |
| Polink-2 Plus AP Rat-NM 18mL Kit / 6mL Kit (no cross react to mouse) | D67-18 / D67-6 | 18mL / 6mL | | | |

Precautions:

Please wear gloves, eye protection and take other necessary precautions. If any of the reagent come in contact with skin wash area completely with plenty of water and soap. If irritation develops seek medical attention.

Remarks:

For research use only.

References:

- De Pasquale A, Paterlini P, Quaglini D. *Immunochemical demonstration of different antigens in single cells in paraffin-embedded histological sections.* Clin Lab Haematol. 1982;4(3):267-72.
- Polak J. M and Van Noorden S. Introduction to Immunocytochemistry Second Edition. Bios Scientific Publishers. P41-54. 1997