

Product datasheet for **AM00003BT-N**

Amyloid beta (1-42 specific) Mouse Monoclonal Antibody [Clone ID: 8G7]

Product data:


Product Type:	Primary Antibodies
Clone Name:	8G7
Applications:	ELISA, IF, WB
Recommended Dilution:	Western blot: 1 µg/ml for HRPO/ECL detection. Recommended blocking buffer: Casein/Tween 20 based blocking and blot incubation buffer. ELISA: 0.05 µg/ml. Immunocytochemistry: 1-10 µg/ml.
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	C-terminal peptide conjugated to KLH
Specificity:	This antibody specifically interacts with the C-Terminus of beta-Amyloid (1-42) and does not crossreact with beta-Amyloid (1-40).
Formulation:	PBS containing 0.09% Sodium Azide as preservative Label: Biotin State: Liquid purified IgG fraction
Concentration:	lot specific
Purification:	Size Exclusion Chromatography
Conjugation:	Biotin
Storage:	Store the antibody (aliquote in liquid nitrogen) at -80°C. Avoid repeated freezing and thawing. Thaw aliquots at 37°C. Thawed aliquots may be stored at 2-8°C up to 3 months.
Stability:	Shelf life: one year from despatch.



[View online »](#)

Background:

The beta-amyloid peptide (beta A4), proteolytically released from the amyloid precursor protein (APP), is the principal component of senile plaques in Alzheimer's disease. Cleavage of APP by alpha-secretase or alternatively by beta-secretase leads to generation and extracellular release of soluble APP peptides, S-APP-alpha and S-APP-beta, respectively, and the retention of corresponding membrane-anchored C-terminal fragments, C83 and C99. Subsequent processing of C83 by gamma-secretase yields P3 peptides. This is the major secretory pathway and is nonamyloidogenic. Alternatively, presenilin/nicastrin-mediated gamma-secretase processing of C99 releases the amyloid beta proteins, amyloid-beta 40 (Abeta40) and amyloid-beta 42 (Abeta42), major components of amyloid plaques, and the cytotoxic C-terminal fragments, gamma-CTF(50), gamma-CTF(57) and gamma-CTF(59).

Product images:

Immunoblot Analysis: Amyloid beta A4 peptides were applied on SDS-PAGE and transferred to a PVDF membrane. The immunoblot was probed with 2g/ml mab bA4 (42)-8G7 for 1h at 15-22C and developed by ECL (exposure time: 30 sec). Lane 1: bA4 (1-40) Lane 2: bA4 (1