

Product datasheet for **AM00006PU-N**

Amyloid beta (free N-term, APP reactive) Mouse Monoclonal Antibody [Clone ID: 19H5]

Product data:

Product Type:	Primary Antibodies
Clone Name:	19H5
Applications:	ELISA, WB
Recommended Dilution:	ELISA: 0.05 µg/ml. Western blot: 1 µg/ml for HRPO/ECL detection. <i>Recommended blocking buffer:</i> Casein/Tween 20 based blocking and blot incubation buffer.
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	N-terminal peptide conjugated to KLH.
Specificity:	This antibody recognizes the <i>free</i> N-Terminus of the βA4 polypeptide. The antibody crossreacts with APP.
Formulation:	1 ml 2 x PBS containing PEG and Sucrose State: Purified State: Lyophilized purified IgG fraction Preservative: 0.09% Sodium Azide
Reconstitution Method:	Restore with 1 ml H ₂ O (15 min, RT).
Purification:	Subsequent Thiophilic Adsorption and Size Exclusion Chromatography
Conjugation:	Unconjugated
Storage:	Prior to reconstitution store at -20°C. Following reconstitution store the antibody undiluted at 2-8°C for one month or (in aliquots) at -80°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.



[View online »](#)

Background:

The beta-amyloid peptide (beta A4), proteolytically released from the amyloid precursor protein (APP), is the principal component of senile plaques in Alzheimers disease. Cleavage of APP by alpha-secretase or alternatively by beta-secretase leads to generation and extracellular release of soluble APP peptides, S-APP-alpha and S-APP-beta, respectively, and the retention of corresponding membrane-anchored C-terminal fragments, C83 and C99. Subsequent processing of C83 by gamma-secretase yields P3 peptides. This is the major secretory pathway and is nonamyloidogenic. Alternatively, presenilin/nicastrin-mediated gamma-secretase processing of C99 releases the amyloid beta proteins, amyloid-beta 40 (Abeta40) and amyloid-beta 42 (Abeta42), major components of amyloid plaques, and the cytotoxic C-terminal fragments, gamma-CTF(50), gamma-CTF(57) and gamma-CTF(59).

Product images:



Immunoblot Analysis: Amyloid beta A4 peptides (lane 1: bA4 (1-40); lane 2: bA4 (1-42); lane 3: bA4 (1-43)) were applied on SDS-PAGE and transferred to a PVDF membrane. The immunoblot was probed with 2g/ml mab bA4N-19H5 for 1h at 15-22C and developed by ECL