

Product datasheet for **AM00019PU-N**

beta Catenin (CTNNB1) (N-term, exon 2) (incl. pos. control) Mouse Monoclonal Antibody [Clone ID: 7D11]

Product data:

Product Type:	Primary Antibodies
Clone Name:	7D11
Applications:	ELISA, IP, WB
Recommended Dilution:	<u>Western Blot</u> : 0.5 µg/ml for HRPO/ECL detection. Recommended blocking buffer: Casein/Tween 20 based blocking and blot incubation buffer. <u>Immunoprecipitation</u> : 1 - 10 µg per 10e6 vanadate treated A431 cells. <u>ELISA</u> : 0.05 µg/ml.
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG2a
Clonality:	Monoclonal
Immunogen:	Recombinant β-Catenin
Specificity:	This antibody specifically interacts with the N-terminus (exon 2) of β-catenin at 90 kDa.
Formulation:	1 ml 2 x PBS / 0.09% Sodium Azide / PEG and Sucrose State: Purified State: Lyophilized purified IgG
Reconstitution Method:	Restore with 1 ml H ₂ O (15 min, RT).
Purification:	Size exclusion chromatography
Conjugation:	Unconjugated
Storage:	Store lyophilized (preferably in a desiccator) at -20°C and reconstituted (aliquote and freeze in liquid nitrogen) at -80°C. Avoid repeated freezing and thawing. Thaw aliquots at 37°C. Thawed aliquots may be stored at 4°C up to 3 months.
Stability:	Shelf life: one year from despatch.
Gene Name:	catenin beta 1
Database Link:	<u>Entrez Gene 1499 Human P35222</u>




[View online »](#)

Background:

The α -, β - and γ -catenins are cytoplasmic proteins mediating the interaction of Ca^{2+} -dependent transmembrane adhesion molecules (cadherins) with the cytoskeletal network. The direct interaction of β -catenin with the cytoplasmic domain of cadherins plays a crucial role for cell-cell adhesion and signal transmission between neighbouring cells. Recent studies indicate that β -catenin may also play a role in tumorigenesis since it forms complexes with the tumor suppressor gene product APC. β -catenin directly interacts and constitutively activates transcription factors of the TCF/LEF gene family. Thus it is proposed that β -catenin plays a dual role not only in the maintenance and regulation of cell-cell interactions but also in the regulation of gene activity.

Synonyms:

CTNNB1, CTNNB, Beta-catenin

Product images:

Detection of endogenous β -Catenin Whole cell lysates of serum starved tumor cells (20,000 cells per lane) were applied to SDS-PAGE and transferred to a PVDF membrane. The immunoblot was probed with mab b-Cat-7D11 (0.5 $\mu\text{g}/\text{ml}$) for 1h at RT and developed by ECL (exp. time: 3 min). lane 1: A431; lane 2:SW480; lane 3: SW620; lane 4: HT29; lane 5: MCF7; lane 6: MDA-MB231; lane 7: T47D