

## Product datasheet for AM00037BT-N

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## EGFR pTyr1092 (incl. pos. control) Mouse Monoclonal Antibody [Clone ID: 15A2]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: 15A2

**Applications:** ELISA, IP, WB

**Recommended Dilution:** Western Blot: 1 µg/ml for HRPO/ECL detection.

Recommended blocking buffer: Casein/Tween 20 based blocking and blot incubation buffer.

ELISA: 0.1 μg/ml (protein ELISA).

Immunoprecipitation: 1 - 10 μg per 106 pervanadate-treated A431 cells.

Luminex.

Reactivity: Canine, Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Phosphopeptide conjugated to KLH.

**Specificity:** This antibody specifically recognizes EGFR phosphorylated at Tyrosine 1092. Mab 15A2 doest

not crossreact with the highly homologoes pTyr 1139 of activated erbB2.

**Formulation:** PBS/0.09% Na-Azide/PEG and Sucrose

Label: Biotin

State: Liquid purified IgG

**Concentration:** lot specific

**Purification:** Size exclusion chromatography

Conjugation: Biotin

Storage: Store the antibody (aliquote in liquid nitrogen) at -80°C.

Avoid repeated freezing and thawing.

Thaw aliquots at 37°C. Thawed aliquots may be stored at 4°C up to 3 months.

**Stability:** Shelf life: one year from despatch.

**Database Link:** Entrez Gene 1956 Human

P00533





## EGFR pTyr1092 (incl. pos. control) Mouse Monoclonal Antibody [Clone ID: 15A2] - AM00037BT-N

Background:

EGFR/erbB receptors are activated upon binding of EGF and EGF-related growth factors such as TGF alpha, beta-cellulin, Hb-EGF, HRG, or NRG. Binding of these ligands leads to receptor homo- and heterodimerization followed by autophosphorylation and activation of downstream signal transduction pathways (MAPK, PI3K/PKB, and STAT). In addition, EGFR becomes fully activated after phosphorylation of Y869 by src family kinases. Phosphorylation of Y1069 leads to association with cbl and subsequent receptor degradation. Phosphorylation of S1071 by CamKinase II leads to attenuation of kinase activity; phosphorylation of T678 (by PKC) and T693 (by MAPK, p38) interferes with receptor endocytosis/recycling.

Synonyms:

Epidermal growth factor receptor, EGF Receptor, erbB-1, c-ErbB-1