

## Product datasheet for **AM00086BT-N**

### **ERK2 (MAPK1) (N-term) (incl. pos. control) Mouse Monoclonal Antibody [Clone ID: 6H3]**

#### Product data:

Product Type:	Primary Antibodies
Clone Name:	6H3
Applications:	ELISA, WB
Recommended Dilution:	<b>ELISA:</b> Use at 0.05 µg/ml. <b>Immunoblotting:</b> 0.5 µg/ml for HRPO/ECL detection. <i>Recommended blocking buffer:</i> Casein/Tween 20 based blocking and blot incubation buffer AS00002BU-N or AS00002BU-L. <i>Included Positive Control:</i> Cell lysate from untreated HepG2 cells (See Protocols for more details).
Reactivity:	Canine, Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Peptide conjugated to KLH <b>Epitope:</b> N-terminus
Specificity:	This antibody specifically recognizes the N-terminus of MAP kinase 2 (ERK2).
Formulation:	PBS with 0.09% Sodium Azide/PEG and Sucrose. Label: Biotin State: Liquid purified IgG fraction.
Concentration:	lot specific
Purification:	Subsequent Thiophilic Adsorption and Size Exclusion Chromatography.
Conjugation:	Biotin
Storage:	Aliquote and freeze in liquid nitrogen Antibody can be stored frozen at -80°C up to 1 year. Thaw aliquots at 37°C. Thawed aliquots may be stored at 4°C up to 3 months.
Gene Name:	mitogen-activated protein kinase 1



[View online »](#)

Database Link:	<a href="#">Entrez Gene 26413 Mouse</a> <a href="#">Entrez Gene 116590 Rat</a> <a href="#">Entrez Gene 5594 Human P28482</a>
Background:	Extracellular signal/mitogen activated protein kinases (erk/MAPK) are a group of proline-directed serine/threonine kinases that are activated by dual phosphorylation of conserved threonine and tyrosine residues within a characteristic T X Y peptide motif. The mitogen-activated kinases erk1 (MAPK1) and erk2 (MAPK2) acquire full enzymatic activity upon phosphorylation of both threonine and tyrosine residues within the sequence motif T E Y.
Synonyms:	Mitogen-activated protein kinase 1, p42-MAPK, ERT1, PRKM1, PRKM2, MAP kinase 2, MAPK2, MAPK1
Note:	<p><b>Mol. weight:</b> 42 kDa</p> <p>Protocol: <b>Positive Control Provided.</b> <b>Cell lysate from untreated HepG2</b></p> <p><b>Description:</b> Cell lysate from untreated HepG2 cells, hepatocellular carcinoma (human)</p> <p><b>Format:</b> Lyophilized cell lysate from serum starved HepG2 cells.</p> <p><b>Reconstitution:</b> Restore by addition of 200 µl H<sub>2</sub>O. After complete solubilization add 200 µl 2x SDS-PAGE sample buffer, mix and incubate at 90°C for 5 min.</p> <p><b>Storage:</b> Aliquote and store frozen. Avoid repeated freeze/thaw cycles.</p> <p><b>Application:</b> The positive control cell lysate is recommended for immunoblot applications. 20 µl of positive control cell lysate correspond to ca. 80.000 cells. Use 20 µl / lane (mini gel) for HRPO/ECL detection of the target proteins.</p> <p><b>Please note:</b> The lyophilized cell lysates contain SDS and are not recommended for applications with native proteins such as immunoprecipitation.</p>
Protein Families:	Druggable Genome, Protein Kinase

**Protein Pathways:**

Acute myeloid leukemia, Adherens junction, Alzheimer's disease, Axon guidance, B cell receptor signaling pathway, Bladder cancer, Chemokine signaling pathway, Chronic myeloid leukemia, Colorectal cancer, Dorso-ventral axis formation, Endometrial cancer, ErbB signaling pathway, Fc epsilon RI signaling pathway, Fc gamma R-mediated phagocytosis, Focal adhesion, Gap junction, Glioma, GnRH signaling pathway, Insulin signaling pathway, Long-term depression, Long-term potentiation, MAPK signaling pathway, Melanogenesis, Melanoma, mTOR signaling pathway, Natural killer cell mediated cytotoxicity, Neurotrophin signaling pathway, NOD-like receptor signaling pathway, Non-small cell lung cancer, Oocyte meiosis, Pancreatic cancer, Pathways in cancer, Prion diseases, Progesterone-mediated oocyte maturation, Prostate cancer, Regulation of actin cytoskeleton, Renal cell carcinoma, T cell receptor signaling pathway, TGF-beta signaling pathway, Thyroid cancer, Toll-like receptor signaling pathway, Type II diabetes mellitus, Vascular smooth muscle contraction, VEGF signaling pathway