

Product datasheet for **AM00702BT-N**

Tfrc Rat Monoclonal Antibody [Clone ID: R17217.1.4]

Product data:

Product Type:	Primary Antibodies
Clone Name:	R17217.1.4
Applications:	FC
Recommended Dilution:	Flow Cytometry.
Reactivity:	Mouse
Host:	Rat
Isotype:	IgG2a
Clonality:	Monoclonal
Specificity:	Anti-mouse CD71 antigen monoclonal antibody reacts with the transferrin receptor (CD71). The expression of CD71 is up-regulated after B cell or T cell activation.
Formulation:	PBS containing 0.09% sodium azide (NaN ₃) as preservative and EIA grade BSA as a stabilizing protein Label: Biotin State: Liquid purified Ig fraction
Concentration:	lot specific
Conjugation:	Biotin
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	transferrin receptor
Database Link:	Entrez Gene 22042 Mouse Q62351
Synonyms:	TfR1, p90, Transferrin receptor protein 1



[View online »](#)

Note:Protocol: **FLOW CYTOMETRY ANALYSIS:****Method:**

1. Prepare a cell suspension in media A. For cell preparations, deplete the red blood cell population with Lympholyte®-M cell separation medium.
2. Wash 2 times.
3. Resuspend the cells to a concentration of 2×10^7 cells/ml in media A. Add 50 μ l of this suspension to each tube (each tube will then contain 1×10^6 cells, representing 1 test).
4. To each tube, add ~ 1.0 μ g of this antibody per 10^6 cells.
5. Vortex the tubes to ensure thorough mixing of antibody and cells.
6. Incubate the tubes for 30 minutes at 4°C.
7. Wash 2 times at 4°C.
8. Add 100 μ l of secondary antibody (Streptavidin-FITC) at a 1/500 dilution.
9. Incubate tubes at 4°C for 30 - 60 minutes (It is recommended that tubes are protected from light since most fluorochromes are light sensitive).
10. Wash 2 times at 4°C.
11. Resuspend the cell pellet in 50 μ l ice cold media B.
12. Transfer to suitable tubes for flow cytometric analysis containing 15 μ l of propidium iodide at 0.5 mg/ml in PBS. This stains dead cells by intercalating in DNA.

Media:

- A. Phosphate buffered saline (pH 7.2) + 5% normal serum of host species + sodium azide (100 μ l of 2M sodium azide in 100 mls).
- B. Phosphate buffered saline (pH 7.2) + 0.5% Bovine serum albumin + sodium azide (100 μ l of 2M sodium azide in 100 mls).

Results:

Tissue Distribution by Flow Cytometry Analysis:
(Representative Histogram)

Mouse Strain: Mouse Hybridomas

Cell Concentration: 1×10^6 cells per test

Antibody Concentration Used: 1.0 μ g/ 10^6 cells

Isotypic Control: Biotin Rat IgG2a