

Product datasheet for AM01322PU-S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD79A (202-216) Mouse Monoclonal Antibody [Clone ID: HM57]

Product data:

Product Type: Primary Antibodies

Clone Name: HM57

Applications: FC, IHC, WB

Recommended Dilution: Western blot.

Immunohistochemistry on Frozen Sections.

Immunohistochemistry on Paraffin Sections: 1/100-1/500. This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH

6.0 is recommended for this purpose.

Recommended Positive Control: Human tonsil.

Flow Cytometry: Use 10 μl of 1/10-1/50 diluted CD79a antibody to label 1x16 cells in 100 μl

(Membrane permeabilisation is required).

Reactivity: Bison, Bovine, Canine, Chicken, Deer, Equine, Guinea Pig, Human, Monkey, Mouse, Porcine,

Rabbit, Rat

Host: Mouse

Isotype: lgG1

Clonality: Monoclonal

Immunogen: Synthetic peptide corresponding to 202-216 amino acid sequence of Human mb-1.

Specificity: This antibody recognizes an epitope within the cytoplasmic domain of CD79a. CD79a, also

known as mb-1, is a 45kD protein that is expressed by B lymphocytes during differentiation

from early pre-B cell stage through to plasma cells.

Formulation: PBS

State: Purified

State: Liquid purified IgG fraction from Tissue Culture Supernatant

Preservative: 0.09% Sodium Azide

Concentration: lot specific

Purification: Affinity Chromatography on Protein G

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.





CD79A (202-216) Mouse Monoclonal Antibody [Clone ID: HM57] - AM01322PU-S

Stability: Shelf life: one year from despatch.

Database Link: Entrez Gene 973 Human

P11912


Background: The CD79a molecule associates with CD79b (B29) to form a heterodimer that is non-

covalently linked to surface immunoglobulin, forming the B-cell receptor (BCR) complex. The CD79a/CD79b heterodimers are also necessary for intracellular signalling following antigen-

binding to surface immunoglobulin.

Synonyms: IGA, MB1, B-Cell marker

Product images:



Staining of paraffin embedded human tonsil with mouse anti human CD79a antibody.