

Product datasheet for AM03039FC-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

DcR2 (TNFRSF10D) Mouse Monoclonal Antibody [Clone ID: TRAIL-R4-01]

Product data:

Product Type: Primary Antibodies

Clone Name: TRAIL-R4-01

Applications: FC

Recommended Dilution: Flow Cytometry: 3 µg/ml.

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: TRAIL-R4 (aa 1-210) - hlgGhc fusion protein

Specificity: The antibody TRAIL-R4-01 reacts with TRAIL-R4, a 42 kDa transmembrane protein expressed

on various blood cells.

Formulation: PBS containing 15 mM Sdium Azide as preservative and 0.2% (w/v) high-grade (Protease free)

BSA as a stabilizing agent.

Label: FITC

State: Liquid purified IgG fraction.

Label: Fluorescein Isothiocyanate Isomer 1

Concentration: lot specific

Conjugation: FITC

Storage: Store the antibody undiluted at 2-8°C.

DO NOT FREEZE!

This product is photosensitive and should be protected from light.

Avoid prolonged exposure to light.

Stability: Shelf life: one year from despatch.

Gene Name: tumor necrosis factor receptor superfamily member 10d

Database Link: Entrez Gene 8793 Human

Q9UBN6





DcR2 (TNFRSF10D) Mouse Monoclonal Antibody [Clone ID: TRAIL-R4-01] - AM03039FC-N

Background: TRAIL-R4 (CD264, TR4, DcR2, TRUNDD), expressed mainly on CD8+ and NK cells, belongs to

receptors of TRAIL, a TNF-like membrane toxic protein that induces apoptosis in many tumour cells, but not in normal cells. TRAIL-R4, however, contains partially truncated death domain, thus it is unable to induce apoptosis and serves as a negative regulator of apoptotic signaling by impairment death-inducing signaling complex (DISC) processing. TRAIL-R4 interacts with death receptor 5 (DR5) in the native DISC in a TRAIL-dependent manner and

prevents its corecruitment with death receptor 4 (DR4).

Synonyms: TNFRSF10D, DCR2, TRAIL-R4, TRUNDD, Decoy receptor 2, TRAIL receptor 4

Protein Families: Druggable Genome, Transmembrane

Protein Pathways: Apoptosis, Cytokine-cytokine receptor interaction, Natural killer cell mediated cytotoxicity