

Product datasheet for AM03058FC-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD36 Mouse Monoclonal Antibody [Clone ID: TR9]

Product data:

Product Type: Primary Antibodies

Clone Name: TR9
Applications: FC

Recommended Dilution: Suitable for Flow Cytometry analysis of human blood cells using 20 μl reagent / 100 μl of

whole blood or 10e6 cells in a suspension.

The content of a vial (2 ml) is sufficient for 100 tests.

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal Immunogen: Platelets.

Specificity: The antibody TR9 reacts with CD36 (GPIIIb), a 85 kDa integral membrane glycoprotein

expressed on platelets, macrophages, endothelial cells, early erythroid cells and

megakaryocytes.

This antibody cross-blocks binding of FITC-labeled standard antibody OKM5.

Anti-CD36 antibodies inhibit adhesive functions (e.g. adherence of infected erythrocytes to

target cells).

Formulation: PBS containing 15 mM sodium azide as preservative and 0.2% (w/v) high-grade BSA (Protease

free) as stabilizer.

Label: FITC

State: Liquid purified Ig fraction.

Label: Conjugated with Fluorescein isothiocyanate under optimum conditions. The reagent is

free of unconjugated and adjusted for direct use

Conjugation: FITC

Storage: Store the antibody in the dark at 2-8°C.

DO NOT FREEZE!

Avoid prolonged exposure to light.

Stability: Shelf life: One year from despatch.

Gene Name: CD36 molecule



CD36 Mouse Monoclonal Antibody [Clone ID: TR9] - AM03058FC-N

Database Link: Entrez Gene 948 Human

P16671


Background: CD36 (fatty acid translocase, FAT) is an 88 kDa ditopic glycosylated protein that belongs to the

class B family of scavenger receptors. CD36 is expressed by most resting marginal zone B cells but not by follicular and B1 B cells, and it is rapidly induced on Follicular B cells in vitro upon TLR and CD40 stimulation. CD36 does not affect the development of B cells, but modulates both primary and secondary antibody response. Similarly to glucose transporter GLUT4, CD36 is translocated from intracellular pools to the plasma membrane following cell stimulation by insulin. In mouse, CD36 is responsible for gustatory perception of long-chain

fatty acids.

Synonyms: Glycoprotein IIIb, PAS IV, PAS-4, Thrombospondin receptor, GP3B, GP4

Product images:



Surface staining of human platelets with anti-CD36 (TR9) FITC.