

Product datasheet for AM03061AC-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD95 (FAS) Mouse Monoclonal Antibody [Clone ID: LT95]

Product data:

Product Type: Primary Antibodies

Clone Name: LT95
Applications: FC

Recommended Dilution: Flow cytometry: 10 μl reagent / 100 μl of whole blood or 10e6 cells in a suspension.

The content of a vial (1 ml) is sufficient for 100 tests.

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: HUT-78 human T cell lymphoma cell line

Specificity: The antibody LT95 reacts with CD95 (Fas/APO-1), a 46 kDa single chain type I glycoprotein of

the tumour necrosis factor/nerve growth factor (TNF/NGF) receptor superfamily, expressed

on a variety of normal and neoplastic cells.

It seems that the antibody LT95 does not induce Fas mediated apoptosis, although it cross-

blocks anti-Fas DX2 antibody that recognizes a functional epitope of Fas molecule.

Formulation: PBS containing 15 mM sodium azide as preservative and 0.2% (w/v) high-grade BSA (Protease

free) as stabilizer

Label: APC

State: Liquid IgG fraction

Label: Conjugated with cross-linked Allophycocyanin under optimum conditions

Purification: Size-exclusion chromatography

Conjugation: APC

Storage: Store the antibody in the dark at 2-8°C.

Do Not Freeze!

Avoid prolonged exposure to light.

Stability: Shelf life: one year from despatch.

Gene Name: Fas cell surface death receptor



Database Link: Entrez Gene 355 Human

P25445


Background: CD95 (Fas, APO-1), a 46 kDa transmembrane glycoprotein, is a cell death receptor of the TNFR

superfamily. Stimulation of CD95 results in aggregation of its intracellular death domains, formation of the death-inducing signaling complex (DISC) and activation of caspases. In type I cells caspase 3 is activated by high amounts of caspase 8 generated at the DISC, in type II cells low concentration of caspase 8 activates pathway leading to the release of cytochrome c from mitochondria and activation of caspase 3 by cytochom c. Besides its roles in induction

of apoptosis, Fas also triggers pro-inflammatory cytokine responses.

Synonyms: FASLG receptor, Apo-1 antigen, APT1, FAS1, TNFRSF6

Product images:



Surface staining of human peripheral blood cells with anti-CD95 (LT95) APC.