

Product datasheet for AM03064PU-N

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436

OriGene Technologies, Inc.

Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product data:

Product Type: Primary Antibodies

Clone Name: LT21
Applications: FC, IHC

Recommended Dilution: Flow cytometry: Recommended dilution: 1-5 µg/ml.

CD21 (CR2) Mouse Monoclonal Antibody [Clone ID: LT21]

Reactivity: Bovine, Canine, Human, Porcine

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: IM9 human B-lymphoblastoid cell line

Specificity: The antibody LT21 reacts with CD21 (CR2), a 145 kDa transmembrane glycoprotein

(complement C3d receptor - C3dR) expressed on B lymphocytes, follicular dendritic cells, some epithelial cells and a subsets of T lymphocytes. It is not expressed on immature B cells.

Formulation: PBS, pH 7.4 containing 15 mM sodium azide as preservative

State: Purified

State: Liquid purified IgG fraction (> 95% pure by SDS-PAGE)

Concentration: lot specific

Purification: Purified by protein-A affinity chromatography.

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C.

DO NOT FREEZE!

Stability: Shelf life: one year from despatch.

Gene Name: complement component 3d receptor 2

Database Link: Entrez Gene 1380 Human

P20023






Background:

CD21 (complement receptor 2, CR2) binds C3 complement fragments, especially its breakdown fragments, which remain covalently attached to complement activating surfaces or antigen. CD21 has important roles in uptake and retention of immunocomplexes, survival of memory B cells and in development and maintenance of the humoral response to T-dependent antigens. CD21 also serves as a key receptor for Epstein-Barr virus binding and is involved in targeting prions to folicular dendritic cells and expediting neuroinvasion following peripheral exposure to prions. A soluble form of the CD21 (sCD21) is shed from the lymphocyte surface and retains its ability to bind respective ligands.

Synonyms:

Complement receptor type 2, CR2, C3DR, C3d receptor, EBV Receptor, Dendritic Cell Marker

Product images:



Double staining of human, canine and porcine B lymphocytes with anti-CD79a (HM57) and anti-CD21 (LT21) antibody.