

## Product datasheet for AM03090PU-N

# OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### CD62L (SELL) Mouse Monoclonal Antibody [Clone ID: LT-TD180]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: LT-TD180

Applications: FC

**Recommended Dilution:** Suitable for **Flow Cytometry**.

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Peripheral blood leukocytes

Specificity: The antibody LT-TD180 reacts with CD62L (L-selectin), a 74-95 kDa single chain type I

glycoprotein expressed on most peripheral blood B lymphocytes, T lymphocytes, monocytes

and granulocytes; it is also present on a subset of NK cells and certain hematopoietic

malignant cells.

**Formulation:** PBS, pH 7.4 with 15 mM sodium azide as preservative

State: Purified

State: Liquid purified IgG fraction (> 95% pure by SDS-PAGE)

**Concentration:** lot specific

**Purification:** Purified by protein-A affinity chromatography.

**Conjugation:** Unconjugated

**Storage:** Store undiluted at 2-8°C.

DO NOT FREEZE!

**Stability:** Shelf life: one year from despatch.

**Gene Name:** selectin L

**Database Link:** Entrez Gene 6402 Human

P14151




#### Background:

CD62L (L-selectin) is an adhesion glycoprotein that is constitutively expressed on the cell surface of leukocytes and mediates their homing to inflammatory sites and peripheral lymph nodes by enabling rolling along the venular wall. CD62L is also involved in activation-induced neutrophil aggregation. Activation-dependent CD62L shedding, however, counteracts neutrophil rolling. CD62L has also signaling roles including enhance of chemokine receptor expression. Similarly to CD62P, the major ligand of CD62L is PSGL-1 (P-selectin glycoprotein ligand-1).

Synonyms:

SELL, LNHR, LYAM1, Leu-8, TQ1, gp90-MEL, LECAM1, LAM-1

## **Product images:**



Surface staining of human peripheral blood cells with anti-CD62L (LT-TD180) purified, GAM-APC.