

Product datasheet for AM03093PU-N

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

CD147 (BSG) Mouse Monoclonal Antibody [Clone ID: MEM-M6/2]

Product data:

Applications:

Product Type: Primary Antibodies

Clone Name: MEM-M6/2

Recommended Dilution: Flow Cytometry (4 µg/ml).

FC. IHC

Immunohistochemistry on Paraffin Sections (10 µg/ml).

Positive Control: Testis tissue.

Reactivity: Human, Porcine

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

Immunogen: A soluble recombinant form of CD147, CD147Rg, which consists of the cDNA coding for the

entire extracellular region of CD147 fused to the DNA coding for the hinge region, CH2 and

CH3 domain of Human IgG1.

Specificity: The antibody MEM-M6/2 recognizes extracellular Ig domain D1 of CD147 (Neurothelin), a 50-

60 kDa type I transmembrane glycoprotein primarily expressed on all leukocytes, red blood

cells, platelets and endothelial cells; it is not expressed by resting lymphocytes.

The antibody MEM-M6/1 is a high-affinity antibody capable of binding to unstimulated

peripheral blood T cells.

Formulation: PBS, pH~7.4

State: Aff - Purified

State: Liquid purified IgG fraction (> 95% pure by SDS-PAGE)

Preservative: 15 mM Sodium Azide

Concentration: lot specific

Purification: Affinity Chromatography on Protein A

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C.

DO NOT FREEZE!

Stability: Shelf life: one year from despatch.





CD147 (BSG) Mouse Monoclonal Antibody [Clone ID: MEM-M6/2] - AM03093PU-N

Gene Name: basigin (Ok blood group)

Database Link: Entrez Gene 682 Human

P35613

Background: CD147 (basigin, neurothelin, OX-47, 5A11, CE9, M6) also known as EMMPRIN (extracellular

matrix metalloproteinase inducer) or TCSF (tumour cell-derived collagenase-stimulatory factor) is an ubiquitously expressed cell surface protein with multiple glycosylated forms. The highest level of CD147 expression is on metabolically active cells, such as lymphoblasts, inflammatory cells, brown adipocytes and malignant tumour cells. CD147 has multiple functions, including facilitating of cell surface expression of monocarboxylate transporter proteins and extracellular matrix metalloproteinases, regulation of integrin functions, it plays

roles in cell development and activation, fetal development or retinal function.

Synonyms: Basigin, BSG, EMMPRIN, TCSF