

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for AM03121LE-N

CD147 (BSG) Mouse Monoclonal Antibody [Clone ID: MEM-M6/6]

Product data:

| Product Type: | Primary Antibodies |
|----------------|--|
| Clone Name: | MEM-M6/6 |
| Reactivity: | Human |
| Host: | Mouse |
| lsotype: | lgG1 |
| Clonality: | Monoclonal |
| Immunogen: | Protein A-CR purified soluble recombinant form of CD147, CD147Rg, which consists of the cDNA coding for the hinge region, CH2-and CH3 domain of human lgG1 (CD147Rg is secreted by transfectants as a dimmer) |
| Specificity: | This antibody recognizes lg domain D2 (membrane proximal) of CD147 (Neurothelin), a 50-60 kDa type l transmembrane glycoprotein primarily expressed on all leukocytes, red blood cells, platelets and endothelial cells; it is not expressed by resting lymphocytes. |
| Formulation: | Azide free phosphate buffered saline (PBS), approx. pH 7.4; 0.2 μm filter sterilized. Endotoxin level is less than 10 EU/mg of the protein, as determined by the LAL test State: Low Endotoxin State: Liquid Ig fraction |
| Concentration: | lot specific |
| Purification: | Protein-A affinity chromatography; > 95% pure (by SDS-PAGE) |
| Conjugation: | Unconjugated |
| Storage: | Store at 2 - 8 °C for up to one month or (in aliquots) at -20 °C for longer. Avoid repeated freezing and thawing. |
| Stability: | Shelf life: one year from despatch. |
| Gene Name: | basigin (Ok blood group) |
| Database Link: | <u>Entrez Gene 682 Human</u> <u>P35613</u> |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

| CD147 (BSG) Mouse Monoclonal Antibody [Clone ID: MEM-M6/6] – AM03121LE-N |
|--|
| CD147 (basigin, neurothelin, OX-47, 5A11, CE9, M6) also known as EMMPRIN (extracellular matrix metalloproteinase inducer) or TCSF (tumour cell-derived collagenase-stimulatory factor) is an ubiquitously expressed cell surface protein with multiple glycosylated forms. The highest level of CD147 expression is on metabolically active cells, such as lymphoblasts, inflammatory cells, brown adipocytes and malignant tumour cells. CD147 has multiple functions, including facilitating of cell surface expression of monocarboxylate transporter proteins and extracellular matrix metalloproteinases, regulation of integrin functions, it plays roles in cell development and activation, fetal development or retinal function. |
| Basigin, BSG, EMMPRIN, TCSF Druggable Genome, ES Cell Differentiation/IPS, Transmembrane |
| |

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US