

Product datasheet for AM03126PU-S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PD1 (PDCD1) (21-167) Mouse Monoclonal Antibody [Clone ID: 4F12]

Product data:

Product Type: Primary Antibodies

Clone Name: 4F12

Applications: ELISA, WB

Recommended Dilution: ELISA.

Western blot: 1/500 (starting dilution) - 1/1000.

Reactivity: Human
Host: Mouse
Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Recombinant human PDCD1 (21-167aa) purified from *E. coli*

Specificity: This antibody detects PDCD1.

Formulation: PBS, pH 7.4 containing 0.02% Sodium Azide and 10% Glycerol

State: Purified

State: Liquid Ig fraction

Concentration: lot specific

Purification: Protein-G affinity chromatography

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for up to two weeks or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: programmed cell death 1

Database Link: Entrez Gene 5133 Human

Q15116






Background:

PDCD1 is an immunoreceptor that belongs to the immunoglobulin (Ig) superfamily, also known as Programmed Cell Death 1. It is expressed on activated T cell, B cells, myeloid cells and a subset of thymocytes. PDCD1 interacts with expressed PDL1 and PDL2 to inhibit the proliferation and activation of T cells and B cells. Both PDL1 and PDL2 are expressed in a wide variety of normal tissues including heart, lung, placenta, spleen and kidney. PDCD1 might be involved in the regulation of peripheral tolerance and may help prevent autoimmune diseases.


Synonyms:

Protein PD-1, hPD-1, PDCD1

Product images:



Western blot analysis Cell lysates of 293T and U937 (40ug) were resolved by SDS-PAGE, transferred to NC membrane and probed with anti-human PDCD1 (1:500). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.



Cell lysates of 293T and U937 (40ug) were resolved by SDS-PAGE, transferred to NC membrane and probed with anti-human PDCD1 (1:500). Proteins were visualized using a goat antimouse secondary antibody conjugated to HRP and an ECL detection system.