

Product datasheet for **AM03182BT-N**

Phosphotyrosine (incl. pos. control) Mouse Monoclonal Antibody [Clone ID: 1F9]

Product data:

Product Type:	Primary Antibodies
Clone Name:	1F9
Applications:	ELISA, IP, WB
Recommended Dilution:	ELISA: Use at 0.05 µg/ml. Immunoblotting: 0.5 µg/ml for HRPO/ECL detection. <i>Recommended blocking buffer:</i> Casein/Tween 20 based blocking and blot incubation buffer AS00002BU-N or AS00002BU-L. Immunoprecipitation: Use at 1-10 µg per 10e6 pervanadate-treated A431 cells. <i>Included Positive Control:</i> Phosphotyrosine MW standard.
Reactivity:	Canine, Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Phosphopeptide conjugated to hemocyanin Epitope: ...P-E-pY-H-N...
Specificity:	Monoclonal antibody 1F9 recognizes Phosphotyrosine in the context of the surrounding amino acids, tolerating charged amino acids directly neighboured to Phosphotyrosine.
Formulation:	PBS with 0.09% Sodium Azide/PEG and Sucrose. Label: Biotin State: Liquid purified IgG fraction.
Concentration:	lot specific
Purification:	Subsequent Thiophilic Adsorption and Size Exclusion Chromatography.
Conjugation:	Biotin
Storage:	Aliquote and freeze in liquid nitrogen. Antibody can be stored frozen at -80°C up to 1 year. Thaw aliquots at 37°C. Thawed aliquots may be stored at 4°C up to 3 months.



[View online »](#)

Background:

Phosphorylation and dephosphorylation of cellular proteins are central steps in transducing extracellular signals to the cell nucleus. Phosphorylated epitopes may serve as docking sites for the assembly of protein complexes or may alter the 3-dimensional protein structure thus modulating enzymatic activity or the ability to undergo protein-protein-interactions. Modification of proteins on tyrosine residues is mediated by protein tyrosine kinases. Tyrosine phosphorylation may alter the biological activity or mediate the assembly of protein complexes via interaction of phosphotyrosine residues with SH2 or PID domains.