

Product datasheet for AM03204FC-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD22 Mouse Monoclonal Antibody [Clone ID: IS7]

Product data:

Product Type: Primary Antibodies

Clone Name: IS7
Applications: FC

Recommended Dilution: Flow Cytometry analysis of human blood cells using 20 μl reagent / 100 μl of whole blood or

10e6 cells in a suspension.

The content of a vial (2 ml) is sufficient for 100 tests.

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Reh human cell line

Specificity: This antibody detects CD22 (BL-CAM), a 130 kDa type I transmembrane glycoprotein

(immunoglobulin superfamily) expressed in the cytoplasm of pro-B and pre-B lymphocytes,

and on the surface of mature and activated B lymphocytes; it is lost on plasma cells,

peripheral blood T lymphocytes, granulocytes and monocytes.

Formulation: Phosphate buffered saline (PBS) containing 15 mM sodium azide and 0.2% (w/v) high-grade

protease free Bovine Serum Albumin (BSA) as a stabilizing agent

Label: FITC

State: Liquid purified Ig fraction

Label: Conjugated with Fluorescein isothiocyanate under optimum conditions. The reagent is

free of unconjugated

Conjugation: FITC

Storage: Store the antibody at 2 - 8 °C. DO NOT FREEZE! This product is photosensitive and should be

protected from light.

Stability: Shelf life: one year from despatch.

Gene Name: CD22 molecule

Database Link: Entrez Gene 933 Human

P20273





CD22 Mouse Monoclonal Antibody [Clone ID: IS7] - AM03204FC-N

Background: CD22, also known as Siglec-2 (sialic acid-binding immunoglobulin-like lectin-2) is a

transmembrane glycoprotein binding alpha2,6-linked sialic acid-bearing ligands. Intracellular domain of CD22 recruits protein tyrosine phosphatase SHP-1 through the immunoreceptor tyrosine-based inhibitory motifs (ITIMs), thus setting a treshold for B cell receptor-mediated activation. CD22 also regulates B-cell response by involvement in controlling the CD19/CD21-Src-family protein tyrosine kinase amplification pathway and CD40 signaling. CD22 exhibits

hallmarks of clathrin-mediated endocytic pathway.

Synonyms: SIGLEC2, Siglec-2, B-cell receptor CD22, Leu-14, BL-CAM