

Product datasheet for AM05270PU-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Alpha SNAP (NAPA) Mouse Monoclonal Antibody [Clone ID: 15D4]

Product data:

Product Type: Primary Antibodies

Clone Name: 15D4

Applications: ELISA, IP, WB

Recommended Dilution: Western Blot: 0.5-5 ug/ml.

Immunoprecipitationon native and denatured samples: 2-10 ug/ml.

Reactivity: Bovine, Human, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Hybridoma produced by the fusion of splenocytes from mice immunized with recombinant

human a-SNAP protein and mouse myeloma cells.

Specificity: This antibody specifically recognizes a-SNAP as a single band of ~36 kDa on western blot of

rat kidney, rat brain and MDBK cells and do not recognize b-SNAP.

Formulation: 20 mM Sodium Phosphate, pH 7.5, 150 mM Sodium Chloride, 50% Glycerol without

preservatives. State: Purified

State: Liquid purified IgG fraction (> 95% pure).

Concentration: lot specific

Purification: Standard chromatographic techniques.

Conjugation: Unconjugated

Storage: Store the antibody at -20°C.

Avoid repeated freezing and thawing.

Stability: Shelf life: One year from despatch.

Gene Name: NSF attachment protein alpha

Database Link: Entrez Gene 140673 RatEntrez Gene 8775 Human

P54920





Alpha SNAP (NAPA) Mouse Monoclonal Antibody [Clone ID: 15D4] - AM05270PU-N

Background:

SNAPs (soluble NSF attachment proteins), acting in concert with SNAREs (SNAP receptors) and the N-ethylmaleimide-sensitive fusion protein (NSF), are required for the fusion of transport vesicles to their target membranes in synaptic transmission, intra-Golgi transport, endosome-to-endosome fusion and transcytotic vesicles-to-plasma membrane transport. Vesicle-to-target membrane docking (initial contact) occurs when the vesicle SNARE binds to its cognate target membrane SNARE. a-SNAP (or bSNAP in brain) then binds to this docking complex and mediates the binding of NSF and thus the formation of a 20 S fusion particle. It is thought that, once NSF is bound, ATP hydrolysis by NSF initiates the fusion process. Whereas a-SNAP is expressed in all mammalian tissues, b-SNAP is expressed only in brain.

Synonyms:

NAPA, SNAP-alpha, SNAPA