

Product datasheet for AM05295PU-N

AKR1C2 Mouse Monoclonal Antibody [Clone ID: T101]

Product data:

Product Type: Primary Antibodies Clone Name: T101 **Applications:** ELISA, IF, IHC, WB Recommended Dilution: Immunohistochemistry. Immunofluorescense. Enzyme Immunoassay. Western blot (1/250-1/500): Detects a 32 kDa band in human breast cancer tissue and Porcine liver tissue. Positive Control: Human breast cancer and Porcine liver tissue. **Reactivity:** Human, Mammalian, Mouse, Porcine Host: Mouse Isotype: lgG1 Monoclonal **Clonality:** Hybridoma produced by the fusion of splenocytes from mice immunized with recombinant Immunogen: type 2 Dihydrodiol Dehydrogenase from Human lung cancer and Mouse myeloma cells. Formulation: PBS containing 0.02% Sodium Azide as preservative. State: Liquid **Conjugation:** Unconjugated Storage: Store the antibody (in aliquots) at -20°C. Avoid repeated freezing and thawing. Stability: Shelf life: One year from despatch. Gene Name: aldo-keto reductase family 1, member C2 Database Link: Entrez Gene 1646 Human P52895



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Serigene AKR1C2 Mouse Monoclonal Antibody [Clone ID: T101] – AM05295PU-N

Background: DDH is a member of aldo-keto reductase superfamily (1,2), which catalyzes reduction of aldehyde or ketone to a corresponding alcohol by using NADH or NADPH as a cofactor. In liver, the enzyme is abundantly located in the cytoplasm as a monomeric 34-36 kDa protein (3,4). Interestingly, by differential display Shen et al. (5) has shown that overexpression of DDH could be identified in ethacrynic acid-induced drug-resistant human colon cancer cells. Detection of DDH overexpression in drug-resistant human stomach cancer cells, which were selected by the gradual adaptation to daunorubicin, further suggested that DDH might be associated with the drug-resistance in cancer cells (6). In a recent study, DDH expression was further shown to have prognostic significance in patients with NSCLC. By using an immunohistochemical method to determine DDH expression in surgical specimens, DDH expression was identified in patients with leukemia, lung cancer, esophageal cancer, transitional cancer and breast cancer. Furthermore, overexpression of DDH was confirmed by immunoblotting and in situ hybridization. Correlation between clinicopathological parameters and DDH expression as well as the prognostic significance of DDH expression in patients with advanced cancer was indicated. 3-alpha-HSD3, Dihydrodiol dehydrogenase 2 Synonyms: **Protein Families:** Druggable Genome

Metabolism of xenobiotics by cytochrome P450

Product images:

Protein Pathways:



Figure 2. Western blot analysis using DDH antibody (AM05295PU-N) on porcine liver (A) and breast cancer (B).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US





Figure 1. Immunoperoxidase staining of formalinfixed paraffin embedded human lung cancer tissue showing cytoplasmic localization of dihydrodiol dehydrogenase.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US