

Product datasheet for AM05570PU-T

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Cd80 Rat Monoclonal Antibody [Clone ID: RM80]

Product data:

Product Type: Primary Antibodies

Clone Name: RM80
Applications: FC, IHC

Recommended Dilution: Flow cytometry: 1/10 - 1/25, use 10 μl of the suggested working dilution to label 1x10e6 cells

in 100 μ l.

Immunohistochemistry on frozen sections.

Functional assays.

Reactivity: Mouse
Host: Rat
Isotype: IgG2a

Clonality: Monoclonal

Immunogen: BCL1 cells expressing CD80

Specificity: This antibody recognises CD80 (B7-1), a 60kD cell surface glycoprotein which is a member of

the CD28/B7 family. In mice, CD80 is expressed on monocytes, peritoneal macrophages and dendritic cells. Expression may be significantly increased upon B lymphocytes by LPS and by

II -4.

Formulation: Phosphate buffered saline, pH 7.4 containing 0,09% sodium azide.

State: Purified

State: Liquid Ig fraction

Concentration: lot specific

Purification: Affinity chromatography on Protein G

Conjugation: Unconjugated

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: CD80 antigen

Database Link: Entrez Gene 12519 Mouse

Q00609





Cd80 Rat Monoclonal Antibody [Clone ID: RM80] - AM05570PU-T

Background:

CD80 is a member of the Ig superfamily and serves as the ligand for two T cell molecules, CD28 and CTLA4. Interactions between CD28 and CD80 on activated B cells result in enhanced T cell activation. CD80 is rapidly induced on the surface of in vitro activated B cells, Epstein Barr Virus (EBV) transformed B cell lines, Burkitts lymphoma cell lines, freshly isolated follicular B lymphoma cells, T cells, and monocytes. It is also expressed at high levels in dendritic cells. It reacts weakly with a small proportion of non activated normal B cells and with HTLV1 infected T cells. CD80 does not react with peripheral monocytes, resting and activated normal T cells, T cell lines and T cell clones, nor with myelomonocytic cell lines.

Synonyms:

CD28LG, CD28LG1, LAB7, BB1, B7.1, B7-1