

Product datasheet for **AM05570RP-N**

Cd80 Rat Monoclonal Antibody [Clone ID: RM80]

Product data:

Product Type:	Primary Antibodies
Clone Name:	RM80
Applications:	FC
Recommended Dilution:	Flow Cytometry: Use 10 ul of neat antibody to label 1x10e6 cells in 100 ul.
Reactivity:	Mouse
Host:	Rat
Isotype:	IgG2a
Clonality:	Monoclonal
Immunogen:	BCL1 cells expressing CD80.
Specificity:	This antibody recognises mouse CD80 (B7-1), a 60kD cell surface glycoprotein which is a member of the CD28/B7 family. In mice, CD80 is expressed on monocytes, peritoneal macrophages and dendritic cells, and expression may be significantly increased upon B lymphocytes by LPS and by IL-4.
Formulation:	PBS, pH 7.4 containing 0.09% Sodium Azide as preservative and 1% BSA + 5% sucrose as stabilizer. Label: PE State: Lyophilized purified IgG fraction. Label: R. Phycoerythrin (RPE)
Reconstitution Method:	Reconstitute with 1.0 ml distilled water.
Concentration:	lot specific
Purification:	Affinity Chromatography on Protein G.
Conjugation:	PE
Storage:	Store the antibody undiluted at 2-8°C. DO NOT FREEZE! This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.
Stability:	Shelf life: One year from despatch.
Gene Name:	CD80 antigen



[View online »](#)

Database Link: [Entrez Gene 12519 Mouse Q00609](#)

Background: CD80 is a member of the Ig superfamily and serves as the ligand for two T cell molecules, CD28 and CTLA4. Interactions between CD28 and CD80 on activated B cells result in enhanced T cell activation. CD80 is rapidly induced on the surface of in vitro activated B cells, Epstein Barr Virus (EBV) transformed B cell lines, Burkitts lymphoma cell lines, freshly isolated follicular B lymphoma cells, T cells, and monocytes. It is also expressed at high levels in dendritic cells. It reacts weakly with a small proportion of non activated normal B cells and with HTLV1 infected T cells. CD80 does not react with peripheral monocytes, resting and activated normal T cells, T cell lines and T cell clones, nor with myelomonocytic cell lines.

Synonyms: CD28LG, CD28LG1, LAB7, BB1, B7.1, B7-1