

Product datasheet for AM05877PU-L

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

Ltbr Rat Monoclonal Antibody [Clone ID: 5G11b]

Product data:

Product Type: Primary Antibodies

Clone Name: 5G11b

Applications: FC

Recommended Dilution: Flow cytometry: use 10 μl of 1/50 - 1/100 dilution to label 10e6 cells in 100 μl.

Reactivity: Mouse

Host: Rat IgG2a

Clonality: Monoclonal

Immunogen: Recombinant extracellular domain of LT beta R

Specificity: This antibody recognises lymphotoxin-beta receptor (LT beta R), a member of the tumour

necrosis factor family. LT beta R is primarily expressed on stromal cells and fibroblasts and at lower levels on myeloid cell lines. Clone 5G11b specifically recognises surface expressed LT beta R. The clone is reported to be agonistic with respect to NF Kappa B activation, MIP-2

induction and IL-6 release (2).

This product is routinely tested on WEH1 cells.

Formulation: Phosphate buffered saline pH7.4, 0.09% Sodium Azide

State: Purified

State: Liquid Ig fraction

Concentration: lot specific

Purification: Affinity chromatography on Protein G

Conjugation: Unconjugated

Storage: Store the antibody at 2 - 8 °C up to one month or (in aliquots) at -20 °C for longer. Avoid

repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: lymphotoxin B receptor

Database Link: Entrez Gene 17000 Mouse

P50284





Ltbr Rat Monoclonal Antibody [Clone ID: 5G11b] - AM05877PU-L

Background:

The protein encoded by this gene is a member of the tumor necrosis factor (TNF) family of receptors. It is expressed on the surface of most cell types, including cells of epithelial and myeloid lineages, but not on T and B lymphocytes. The protein specifically binds the lymphotoxin membrane form (a complex of lymphotoxin alpha and lymphtoxin beta). The encoded protein and its ligand play a role in the development and organization of lymphoid tissue and tranformed cells. Activation of the encoded protein can trigger apoptosis.

Synonyms:

Tumor necrosis factor receptor 3, TNF-R3, Tumor necrosis factor C receptor, TNFCR, TNFRSF3, D12S370, Lymphotoxin-beta receptor