

Product datasheet for AM05877RP-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

LTBR Rat Monoclonal Antibody [Clone ID: 5G11b]

Product data:

Product Type: Primary Antibodies

Clone Name: 5G11b

Applications: FC

Recommended Dilution: Flow Cytometry: Use 10 µl of the suggested working dilution to label 10e6 cells in 100 µl.

Reactivity: Mouse

Host: Rat IgG2a

Clonality: Monoclonal

Immunogen: Recombinant extracellular domain of LT beta R

Specificity: This antibody recognises lymphotoxin-beta receptor (LT beta R), a member of the tumour

necrosis factor family. LT beta R is primarily expressed on stromal cells and fibroblasts and at lower levels on myeloid cell lines. Clone 5G11b specifically recognises surface expressed LT beta R. The clone is reported to be agonistic with respect to NF Kappa B activation, MIP-2

induction and IL-6 release (2).

This product is routinely tested on mouse WEHI cells.

Formulation: Phosphate buffered saline pH7.4, 0.09% Sodium Azide, 1% Bovine Serum Albumin

Label: PE

State: Lyophilized Ig fraction Label: R. Phycoerythrin (RPE)

Reconstitution Method: Reconstitute with 1.0 ml distilled water

Concentration: lot specific

Purification: Affinity chromatography on Protein G

Conjugation: PE

Storage: Store the antibody prior and following reconstitution at 2-8°C.

DO NOT FREEZE.

This product is photosensitive and should be protected from light.

Stability: Shelf life: One year from despatch.

Gene Name: lymphotoxin beta receptor





LTBR Rat Monoclonal Antibody [Clone ID: 5G11b] - AM05877RP-N

Database Link: P36941

Background: The protein encoded by this gene is a member of the tumor necrosis factor (TNF) family of

receptors. It is expressed on the surface of most cell types, including cells of epithelial and myeloid lineages, but not on T and B lymphocytes. The protein specifically binds the lymphotoxin membrane form (a complex of lymphotoxin alpha and lymphtoxin beta). The encoded protein and its ligand play a role in the development and organization of lymphoid

tissue and tranformed cells. Activation of the encoded protein can trigger apoptosis.

Synonyms: Tumor necrosis factor receptor 3, TNF-R3, Tumor necrosis factor C receptor, TNFCR, TNFRSF3,

D12S370, Lymphotoxin-beta receptor