

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for AM06578SU-N

MEK6 (MAP2K6) Mouse Monoclonal Antibody [Clone ID: 3H12B9]

Product data:

Product Type:	Primary Antibodies
Clone Name:	3H12B9
Applications:	ELISA, IHC, WB
Recommended Dilution:	ELISA: 1/10000. Western Blot: 1/500 - 1/2000. Immunohistochemistry on Paraffin Sections: 1/200 - 1/1000.
Reactivity:	Human, Mouse
Host:	Mouse
lsotype:	lgG1
Clonality:	Monoclonal
Immunogen:	Purified recombinant fragment of Human MAP2K6 expressed in E. Coli.
Specificity:	Recognizes MAP2K6
Formulation:	State: Ascites State: Ascitic fluid containing 0.03% Sodium Azide.
Conjugation:	Unconjugated
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	38 kDa
Gene Name:	mitogen-activated protein kinase kinase 6
Database Link:	<u>Entrez Gene 5608 Human</u> <u>P52564</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


Serigene MEK6 (MAP2K6) Mouse Monoclonal Antibody [Clone ID: 3H12B9] – AM06578SU-N

Background:This gene encodes a member of the dual specificity protein kinase family, which functions as
a mitogen-activated protein (MAP) kinase kinase. MAP kinases, also known as extracellular
signal-regulated kinases (ERKs), act as an integration point for multiple biochemical signals.
This protein phosphorylates and activates p38 MAP kinase in response to inflammatory
cytokines or environmental stress. As an essential component of p38 MAP kinase mediated
signal transduction pathway, this gene is involved in many cellular processes such as stress
induced cell cycle arrest, transcription activation and apoptosis. Tissue specificity: Isoform 2 is
only expressed in skeletal muscle. Isoform 1, on the other hand, is found in skeletal muscle,
heart, and in lesser extent in liver or pancreas.


Synonyms:

MAPKK 6, MAPK/ERK kinase 6, SAPKK3, MEK6, MKK6, MAP kinase kinase 6


Product images:



Western blot analysis using MAP2K6 antibody Cat.-No AM06578SU-N against HepG2 (1), MCF-7 (2) and NIH/3T3 (3) cell lysate.



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Immunohistochemical analysis of paraffinembedded ovarian cancer (left) and kidney cancer (right) using MAP2K6 antibody Cat.-No AM06578SU-N with DAB staining.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US