

Product datasheet for AM08023AC7-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Cd8a Rat Monoclonal Antibody [Clone ID: 53-6.7]

Product data:

Product Type: Primary Antibodies

Clone Name: 53-6.7

Applications: FC, IHC, IP

Recommended Dilution: Flow Cytometry: < / = 0.2 µg/10e6 cells. (Ref.8,9)

Immunohistochemistry (Acetone-Fixed Frozen Sections). (See Ref. 10)

Immunoprecipitation. (Ref.8)

Fractionation of CD8a+ T cells. (Ref.10)

Reactivity: Mouse
Host: Rat
Isotype: IgG2a

Clonality: Monoclonal

Specificity: This antibody recognises the alpha chain of the Murine CD8 heterodimer (Mr. 32-34 kDa).

Formulation: PBS containing 0.09% Sodium Azide as preservative and a stabilizing agent.

Label: APC-Cy7

State: Liquid purified Ig fraction. Label: Allophycocyanin-Cyanine 7

Concentration: lot specific **Conjugation:** APC-Cy7

Storage: Store the antibody undiluted at 2-8°C.

DO NOT FREEZE!

This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Database Link: Entrez Gene 12525 Mouse

P01731





Cd8a Rat Monoclonal Antibody [Clone ID: 53-6.7] - AM08023AC7-N

Background:

In the mouse, CD8 exists in two forms: (i) a CD8 heterodimer composed of an alpha chain (CD8a/Lyt-2) and a beta chain (CD8b/Lyt-3); and (ii) a homodimer of two alpha chains. The heterodimer is found on the surface of essentially all thymocytes and the "suppressor/cytotoxic" subpopulation of mature T lymphocytes. Subsets of intestinal intraepithelial lymphocytes express CD8a without CD8b. It has been suggested that CD8+b-T cells mature extrathymically, while development of the CD8a+b+ population of T cells is thymus-dependent. CD8 acts as a coreceptor with MHC Class I-restricted T cell receptors in antigen recognition and positive selection of MHC class I-restricted CD8+ T cells. In vivo and in vitro treatment with the 53-6.7 monoclonal antibody effectively depletes CD8a+cells. The 53-6.7 monoclonal antibody also blocks allogeneic help specific for class I MHC antigens and T cell responses to IL-2. (Ref. 1-8)

Synonyms:

CD8 alpha chain, CD8A, MAL