

Product datasheet for **AM08023PU-N**

Cd8a Rat Monoclonal Antibody [Clone ID: 53-6.7]

Product data:

Product Type:	Primary Antibodies
Clone Name:	53-6.7
Applications:	FC, FN, IHC, IP
Recommended Dilution:	Flow Cytometry: $< / = 2 \mu\text{g}/10\text{e6 cells}$. (Ref.8,9) Immunohistochemistry (Acetone-Fixed Frozen Sections) (See Ref. 10) Immunoprecipitation. (Ref.8) <i>In vivo</i> and <i>in vitro</i> depletion of CD8a+ cells. (Ref. 1,4,5) Fractionation of CD8a+ T cells. (Ref.10)
Reactivity:	Mouse
Host:	Rat
Isotype:	IgG2a
Clonality:	Monoclonal
Specificity:	This antibody recognises the alpha chain of the Murine CD8 heterodimer (Mr. 32-34 kDa). In vivo and in vitro treatment with the 53-6.7 monoclonal antibody effectively depletes CD8a+ cells. The 53-6.7 monoclonal antibody also blocks allogeneic help specific for class I MHC antigens and T cell responses to IL-2. (Ref. 1-8)
Formulation:	100 mM Borate Buffered Saline, pH 8.2. No preservatives or amine-containing buffer salts added. State: Purified State: Liquid purified Ig fraction.
Concentration:	lot specific
Conjugation:	Unconjugated
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	CD8 antigen, alpha chain
Database Link:	Entrez Gene 12525 Mouse P01731



[View online »](#)

Background:

In the mouse, CD8 exists in two forms: (i) a CD8 heterodimer composed of an alpha chain (CD8a/Lyt-2) and a beta chain (CD8b/Lyt-3); and (ii) a homodimer of two alpha chains. The heterodimer is found on the surface of essentially all thymocytes and the "suppressor/cytotoxic" subpopulation of mature T lymphocytes. Subsets of intestinal intraepithelial lymphocytes express CD8a without CD8b. It has been suggested that CD8^b- T cells mature extrathymically, while development of the CD8^a+^b+ population of T cells is thymus-dependent. CD8 acts as a coreceptor with MHC Class I-restricted T cell receptors in antigen recognition and positive selection of MHC class I-restricted CD8⁺ T cells.

Synonyms:

CD8 alpha chain, CD8A, MAL