

## Product datasheet for AM08024AC-N

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Cd5 Rat Monoclonal Antibody [Clone ID: 4H8E6]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: 4H8E6

**Applications:** FC, FN, IHC, IP, WB

**Recommended Dilution:** Flow Cytometry: Identification and enumeration of CD5+ cells (< / = 0.2 µg/10e6 cells).

(Ref.15)

**Immunoprecipitation.** (Ref.15)

Immunohistochemistry on Frozen Sections. (Ref.15)

T Cell Activation. (Ref.15) Western blotting. (Ref.15)

Reactivity: Mouse

Host: Rat

**Isotype:** lgG2a

Clonality: Monoclonal

**Specificity:** This antibody is specific to CD5/Lyt-1 (Mr. 67 kDa)

**Formulation:** PBS containing 0.09% Sodium Azide as preservative and a stabilizing agent.

Label: APC

State: Liquid purified Ig fraction.

Label: Allophycocyanin

**Concentration:** lot specific

Conjugation: APC

**Storage:** Store the antibody undiluted at 2-8°C.

**DO NOT FREEZE!** 

This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

Gene Name: CD5 antigen

Database Link: Entrez Gene 12507 Mouse

P13379





## Cd5 Rat Monoclonal Antibody [Clone ID: 4H8E6] - AM08024AC-N

Background:

CD5/Lyt-1 antigen is a monomeric type I transmembrane glycoprotein expressed on thymocytes, T lymphocytes, and a subset of B lymphocytes, but not on natural killer (NK) cells. (Ref.1-3) It has been identified as the major ligand of the B-cell antigen CD72. (Ref.4,5) The frequency of CD5+ B cells exhibits strain-dependent variation, and the phenotypic, anatomical, functional, developmental, and pathological characteristics of the CD5+ B cells suggest that they may represent a distinct lineage, known as B-1 cells (reviewed in Ref.6). Binding of CD5 on the T cell surface can augment alloantigen- or mitogen-induced lymphocyte proliferation and induces increased cytosolic free calcium, IL-2 secretion, and IL-2R expression. (Ref.7-12) It has been proposed that CD5 negatively regulates signal transduction mediated by the T-cell and B-cell receptors. (Ref.13,14)

Synonyms:

CD5, LEU1