

## Product datasheet for AM08024FC-S

## Cd5 Rat Monoclonal Antibody [Clone ID: 4H8E6]

## **Product data:**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Clone Name:	4H8E6
Applications:	FC, FN, IHC, IP, WB
Recommended Dilution:	Flow Cytometry: Identification and enumeration of CD5+ cells (< / = 1 μg/10e6 cells). (Ref.15) Immunoprecipitation. (Ref.15) Immunohistochemistry on Frozen Sections. (Ref.15) T Cell Activation. (Ref.15) Western blotting. (Ref.15)
Reactivity:	Mouse
Host:	Rat
lsotype:	lgG2a
Clonality:	Monoclonal
Specificity:	This antibody is specific to CD5/Lyt-1 (Mr. 67 kDa)
Formulation:	PBS containing 0.09% Sodium Azide as preservative. Label: FITC State: Liquid purified Ig fraction. Label: Fluorescein Isothiocyanate Isomer 1
Concentration:	lot specific
Conjugation:	FITC
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	CD5 antigen
Database Link:	<u>Entrez Gene 12507 Mouse</u> <u>P13379</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Cd5 Rat Monoclonal Antibody [Clone ID: 4H8E6] – AM08024FC-S
Background:	CD5/Lyt-1 antigen is a monomeric type I transmembrane glycoprotein expressed on thymocytes, T lymphocytes, and a subset of B lymphocytes, but not on natural killer (NK) cells. (Ref.1-3) It has been identified as the major ligand of the B-cell antigen CD72. (Ref.4,5) The frequency of CD5+ B cells exhibits strain-dependent variation, and the phenotypic, anatomical, functional, developmental, and pathological characteristics of the CD5+ B cells suggest that they may represent a distinct lineage, known as B-1 cells (reviewed in Ref.6). Binding of CD5 on the T cell surface can augment alloantigen- or mitogen-induced lymphocyte proliferation and induces increased cytosolic free calcium, IL-2 secretion, and IL- 2R expression. (Ref.7-12) It has been proposed that CD5 negatively regulates signal transduction mediated by the T-cell and B-cell receptors. (Ref.13,14)
Synonyms:	CD5, LEU1

~\_\_\_\_\_

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US