

# Product datasheet for AM08026C5-N

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Cd3e Rat Monoclonal Antibody [Clone ID: C363.29B (YCD3-1)]

### **Product data:**

Product Type: Primary Antibodies
Clone Name: C363.29B (YCD3-1)

**Applications:** FC

**Recommended Dilution:** Flow Cytometry: Identification and enumeration of CD3+ (<  $/ = 1 \mu g/10e6$  cells). (Ref.1,2).

Reactivity: Mouse
Host: Rat
Isotype: IgG2b

Clonality: Monoclonal

**Specificity:** This antibody is specific to CD3e (Mr. 25 kD).

**Formulation:** PBS containing 0.09% Sodium Azide as preservative.

Label: Cy5

State: Liquid purified Ig fraction.

Label: Cyanine 5

**Concentration:** lot specific

**Conjugation:** Cy5

**Storage:** Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: CD3 antigen, epsilon polypeptide

Database Link: Entrez Gene 12501 Mouse

P22646





### Cd3e Rat Monoclonal Antibody [Clone ID: C363.29B (YCD3-1)] - AM08026C5-N

Background:

CD3e, a member of the immunoglobulin superfamily of cell surface receptors, is comprised of five invariable chain ranging in size from 16-28 kDa and is closely associated with the T cell antigen receptor (TCR). It is expressed on all T cells of all mouse strains. (Ref.1,2) CD3 plays a major role in signaling during antigen recognition, leading to T-cell activation. The monoclonal antibody C363.29B recognizes an epitope on the 25kD epsilon chain of the CD3/TCR complex. (Ref.1) In the presence of Fc receptor-bearing accessory cells, soluble C363.29B can activate primed and naïve T cell in vitro. Immobilized C363.29 monoclonal antibody can also activate both normal T lymphocytes and cloned T cell lines provided the appropriate accessory signals are present. (Ref.1,3,4) The monoclonal antibody is cytolytic, easily used for cell surface staining and a good

immunoprecipitating antibody. (Ref.1)

Synonyms: T3/Leu-4