

Product datasheet for AM08032BT-N

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

Cd19 Mouse Monoclonal Antibody [Clone ID: MB19-1]

Product data:

Product Type: Primary Antibodies

Clone Name: MB19-1
Applications: FC, IP

Recommended Dilution: Flow Cytometry: < / = 1 μ g/10e6 cells. (Ref.1-3)

Immunoprecipitation. (Ref.1)

Reactivity: Mouse
Host: Mouse
Isotype: IgA

Clonality: Monoclonal

Specificity: This antibody is specific to CD19 B cell differentiation antigen (Mr. 95 kDa).

Formulation: PBS containing Sodium Azide as preservative.

Label: Biotin

State: Liquid purified Ig fraction.

Concentration: lot specific
Conjugation: Biotin

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: CD19 antigen

Database Link: Entrez Gene 12478 Mouse

P25918



Cd19 Mouse Monoclonal Antibody [Clone ID: MB19-1] - AM08032BT-N

Background:

CD19 is a monomeric transmembrane glycoprotein expressed at relatively constant levels throughout B cell development from early pro-B/pre-B cells (i.e., B220+/CD43+/HSA+) through fully differentiated B cell stages. (Ref.1-7) Terminally differentiated plasma cells do not express CD19.1 In humans, the CD19 molecule on the surface of mature B cells associates with CD21 (CR-2) and CD81 (TAPA-1), and this multimolecular complex synergizes with surface immunoglobulin to provide signal transduction and promote cellular activation. (Ref.3,4) All splenic and peritoneal IgM+ cells of both B-1 and B-2 lineages are CD19+, with B-1 cells expressing higher levels of CD19 than B-2 cells in these sites. (Ref.1,7) Recent studies with CD19-deficient mice have suggested that this molecule may not be required for normal generation and maturation of B cells in the bone marrow. (Ref.7)

Synonyms:

Leu-12, B-cell marker