

Product datasheet for AM08033LE-N

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

Cd22 Rat Monoclonal Antibody [Clone ID: 2D6]

Product data:

Product Type: Primary Antibodies

Clone Name: 2D6
Applications: FC, FN

Recommended Dilution: Flow Cytometry.(Ref.2)

Functional Assay: Blocking of B cell homotypic adhesion. (Ref.2)

Reactivity: Mouse

Host: Rat lgG1

Clonality: Monoclonal

Specificity: This antibody is specific to CD22 (Mr 150 kDa), a B Lymphocyte Adhesion Molecule.

Formulation: PBS without preservatives.

State: Low Endotoxin

State: Liquid purified Ig fraction.

Concentration: lot specific

Conjugation: Unconjugated

Storage: Store the antibody undiluted (in aliquots) at -20°C.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: CD22 antigen

Database Link: Entrez Gene 12483 Mouse

P35329



Cd22 Rat Monoclonal Antibody [Clone ID: 2D6] - AM08033LE-N

Background:

CD22, also known as Lyb-8, is a heterodimeric transmembrane glycoprotein and a member of a structurally related group of immunoglobulin (Ig) superfamily domain-containing proteins called the sialoadhesin family. It is detected in the cytoplasm early in B cell development (late pro-B cell stage), appears on the cell surface concomitant with IgD, and is found on most IgM+IgD+ mature B lymphocytes. Expression is lost with terminal differentiation of B cells and is absent on plasma cells. Activation of B cells via cross-linking of surface Ig increases CD22 expression. (Ref.1,2) CD22 associates with the B cell receptor (BCR) complex and mediates intercellular adhesion. (Ref.2-4) Its intracellular domain is phosphorylated after antigen receptor crosslinking and is involved in negative regulation of B-cell activation. (Ref.6-9)

Synonyms:

SIGLEC2, Siglec-2, B-cell receptor CD22, Leu-14, BL-CAM