

## Product datasheet for AM08038LE-N

## 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

## Cd28 Hamster Monoclonal Antibody [Clone ID: PV-1]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: PV-1

Applications: FC, FN, IP

Recommended Dilution: Flow Cytometry. (Ref. 8-10)

Immunoprecipitation. (Ref.8)

Functional Assay: In vitro costimulation of T cells and NK cells. (Ref.8,9)

**Reactivity:** Mouse

**Host:** Hamster

**Isotype:** lgG

Clonality: Monoclonal

**Specificity:** This antibody is specific to Murine CD28.

**Formulation:** PBS containing no preservatives.

State: Low Endotoxin

State: Liquid purified Ig fraction.

**Concentration:** lot specific

**Conjugation:** Unconjugated

**Storage:** Store the antibody undiluted (in aliquots) at -20°C.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

Gene Name: CD28 antigen

**Database Link:** Entrez Gene 12487 Mouse

P31041



## Cd28 Hamster Monoclonal Antibody [Clone ID: PV-1] - AM08038LE-N

**Background:** CD28 is a type I disulfide-linked homodimer that is constitutively expressed on most

thymocytes, at low density on nearly all CD4+ and CD8+ peripheral T lymphocytes, and at very low levels on NK cells. Its expression is upregulated upon T-cell activation. (Ref.1-4) CD28 is a ligand for CD80/B7-1 and CD86/B7-2 on B cells and other antigen presenting cells, and plays an important role in the interaction between T cells and B cells. (Ref.5) CD28 is a co-stimulatory receptor involved in many, but not all, T-cell independent immune responses.

(Ref.6-9)

Synonyms: TP44