

Product datasheet for AM08047LE-N

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

CD49e / ITGA5 Rat Monoclonal Antibody [Clone ID: 5H10-27]

Product data:

Product Type: Primary Antibodies

Clone Name: 5H10-27
Applications: FC, FN, IHC

Recommended Dilution: Flow Cytometry. (Ref.1)

Immunohistochemistry on Frozen Sections: (Ref.7)

Functional Assays:

In vitro blocking of adhesion. (Ref.1,2-4,6)

T-cell co-stimulation. (Ref.3)

Reactivity: Mouse

Host: Rat IgG2a

Clonality: Monoclonal

Immunogen: C57BL/6 x A/J F1 mouse mast cell line MC/9. (Ref.1)

Specificity: This antibody is specific to alpha-5 subunit of the integrin alpha-5 beta-1 Fibronectin

Receptor.

It reacts with the alpha-5 chain (VLA-5) of the CD49e/CD29 heterodimeric Fibronectin

Receptor. (Ref.1-5)

Soluble 5H10-27 antibody inhibits VLA-5-mediated functions in vitro. (Ref.1,2-4,6) Immobilized 5H10-27 has also been demonstrated to co-stimulate the proliferative response of CD8+ T

cells to plate-bound anti-CD3 monoclonal antibody. (Ref.3)

Formulation: PBS containing no preservatives.

State: Low Endotoxin

State: Liquid purified Ig fraction.

Concentration: lot specific

Conjugation: Unconjugated

Storage: Store the antibody undiluted (in aliquots) at -20°C.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: integrin alpha 5 (fibronectin receptor alpha)





CD49e / ITGA5 Rat Monoclonal Antibody [Clone ID: 5H10-27] – AM08047LE-N

Database Link: Entrez Gene 16402 Mouse

P11688

Background: Integrin alpha 5 belongs to the integrin alpha chain family. Integrins are heterodimeric

integral membrane proteins composed of an alpha chain and a beta chain. This gene

encodes the integrin alpha 5 chain. Alpha chain 5 undergoes post translational cleavage in the extracellular domain to yield disulfide linked light and heavy chains that join with beta 1 to form a fibronectin receptor. In addition to adhesion, integrins are known to participate in cell surface mediated signalling. It is expressed on thymocytes, activated T cells, mast cells, and a

variety of mouse cell lines.

Synonyms: Integrin alpha-5, Fibronectin receptor subunit alpha, Integrin alpha-F, VLA-5, VLA5, FNRA