

Product datasheet for AM08052RP-S

Cd80 Rat Monoclonal Antibody [Clone ID: 1G10]

Product data:

Product Type: Primary Antibodies

Clone Name: 1G10 Applications: FC

Recommended Dilution: Flow Cytometry: < / = 0.2 µg/10e6 cells. (Ref.2,7)

Reactivity: Mouse
Host: Rat
Isotype: IgG2a

Clonality: Monoclonal

Specificity: This antibody recognises CD80/B7-1 (Mr. 45-47 kDa)

Formulation: PBS containing 0.09% Sodium Azide as preservative and a stabilizing agent.

Label: PE

State: Liquid purified Ig fraction.

Label: R-Phycoerythrin

Concentration: lot specific

Conjugation: PE

Storage: Store the antibody undiluted at 2-8°C.

DO NOT FREEZE!

This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: CD80 antigen

Database Link: Entrez Gene 12519 Mouse

Q00609



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Cd80 Rat Monoclonal Antibody [Clone ID: 1G10] - AM08052RP-S

Background:

CD80 (B7-1) is a type I transmembrane glycoprotein and a member of the immunoglobulin superfamily of cell surface receptors. It is constitutively expressed by dendritic cells, monocytes, and peritoneal macrophages. Expression of CD80 on B cells is upregulated after activation by various means including stimulation by LPS, IL-4, and surface Ig receptor cross linking with peak expression occurring between 48 and 72 hours postactivation. (Ref.1-4) The interaction of CD80/B7-1 with its ligands, CD28 and CD152/CTLA-4, provides a potent costimulatory signal for T cells activated through the CD3/T cell receptor complex. (Ref.5,6). This Monoclonal antibody 1G10 blocks binding of CTLA-4 to CD80, but it does not block stimulation of T cells by natural antigenpresenting cells. (Ref.2,7).

Synonyms:

CD28LG, CD28LG1, LAB7, BB1, B7.1, B7-1