

Product datasheet for AM08053BT-N

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

Cd86 Rat Monoclonal Antibody [Clone ID: GL1]

Product data:

Product Type: Primary Antibodies

Clone Name: GL1
Applications: FC

Recommended Dilution: Flow Cytometry: < / = 1 μ g/10e6 cells. (Ref.1-7)

Reactivity: Mouse
Host: Rat
Isotype: IgG2a

Clonality: Monoclonal

Immunogen: LPS-activated mouse B cells. (Ref.1)

Specificity: This antibody recognises CD86, the B7-2 co-stimulatory molecule and a ligand for CD28 and

CD152/CTLA-4.

Formulation: PBS containing 0.09% Sodium Azide as preservative.

Label: Biotin

State: Liquid purified Ig fraction.

Concentration: lot specific
Conjugation: Biotin

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: CD86 antigen

Database Link: Entrez Gene 12524 Mouse

P42082





Cd86 Rat Monoclonal Antibody [Clone ID: GL1] - AM08053BT-N

Background:

CD86 (also known as B7-2) is a type I transmembrane glycoprotein and a member of the immunoglobulin superfamily of cell surface receptors. It is expressed at high levels on resting peripheral monocytes and dendritic cells and at very low density on resting B and T lymphocytes. (Ref.1-5) CD86 expression is rapidly upregulated by B-cell specific stimuli with peak expression at 18-42 hours after stimulation. (Ref.1,5) CD86, along with CD80/B7-1, is an important accessory molecule in T cell costimulation via its interaction with CD28 and CD152/CTLA-4. Since CD86 has rapid kinetics of induction, it is believed to be the major CD28 ligand expressed early in the immune response. (Ref.1-9). The monoclonal antibody GL1 blocks mixed lymphocyte reactions and stimulation of T cells by antigen-presenting cells. (Ref.1,3,5)

Synonyms:

CD28LG2, Activation B7-2 antigen, B70, BU63, B7.2, FUN-1