

Product datasheet for AM08054FC-S

Thy1 Rat Monoclonal Antibody [Clone ID: G7]

Product data:

Product Type: Primary Antibodies

Clone Name: G7 FC **Applications:**

Recommended Dilution: Flow Cytometry: $\leq 1 \mu g/10^6$ cells. (Ref.1, 6-8)

Reactivity: Mouse Host: Rat Isotype: lgG2c

Monoclonal Clonality:

Immunogen: Mouse T cell hybridoma C6/G8

Specificity: This antibody recognises Thy-1.1 and Thy-1.2 alloantigens (Thy-1 epitope region A), Mr 28-32

kDa.

Formulation: PBS containing 0.09% Sodium Azide as preservative.

Label: FITC

State: Liquid purified Ig fraction.

Label: Fluorescein Isothiocyanate Isomer 1

Concentration: lot specific

FITC Conjugation:

Storage: Store the antibody undiluted at 2-8°C.

This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: thymus cell antigen 1, theta Database Link: Entrez Gene 21838 Mouse

P01831



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

CD90/Thy-1, a GPI-anchored molecule and one of the smallest members of the immunoglobulin superfamily of cell surface receptors, consists of a single V-set domain. (Ref.1-4) It is expressed on thymocytes, peripheral T lymphocytes, some intraepithelial T lymphocytes and neurons of all mouse strains. (Ref.1,2) MAb G7 stimulates T-cell proliferation and IL-2 secretion, via signalling through the T-cell receptor/CD3 complex. (Ref.1,5-7). It has also been reported to promote apoptosis of thymocytes and CTL clones (Ref.4,7) and to mediate adhesion of thymocytes to thymic stroma. (Ref.8)

Synonyms: Thy-1, THY1, CDw90

Product images:



Immunofluorescent Staining: BALB/c mesenteric lymph node cells were stained with either Rat IgG2c-FITC (as a Negative Control) or Rat anti-Mouse CD90-FITC. Lymphocytes were then gated and analyzed on a FACScan (TM) flow cytometer (BDIS, San Jose, CA). Amount Used: 1 ug/10e6 cells.