

Product datasheet for AM08062PU-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD45 / LCA (CD45.1 alloantigen) Mouse Monoclonal Antibody [Clone ID: A20]

Product data:

Product Type: Primary Antibodies

Clone Name: A20

Applications: FC, FN, IF, IP

Recommended Dilution: Flow Cytometry and Fluorescence Microscopy. (Ref. 3,4)

Immunoprecipitation.

Markers for origin of donor and host cells. (Ref. 3,4)

Reactivity: Mouse
Host: Mouse
Isotype: IgG2a

Clonality: Monoclonal

Specificity: The monoclonal antibody A20 recognizes all mouse leukocytes expressing the CD45.1

allotype.

Formulation: 100 mM Borate Buffered Saline, pH 8.2.

No preservatives or amine-containing buffer salts added.

State: Purified

State: Liquid purified Ig fraction.

Concentration: lot specific

Conjugation: Unconjugated

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: protein tyrosine phosphatase, receptor type, C

Database Link: Entrez Gene 19264 Mouse

P06800



CD45 / LCA (CD45.1 alloantigen) Mouse Monoclonal Antibody [Clone ID: A20] - AM08062PU-N

Background:

The CD45 antigen is an essential regulator of leukocyte activation and development, and is now known to be a protein tyrosine phosphatase, e.g. CD45 is able to dephosphorylate the Src family of protein kinases. (Ref.1,2)

This activity is now known to be required for signal transduction induced by T and B cell antigen receptor engagement, cytokine signaling and Fc receptor stimulation. (Ref.1,2) The functional state of CD45 also has a pronounced effect on lymphocyte development. CD45 exists in multiple forms, such as the restricted variants (CD45R; see companion products AM08044, AM08045, AM08046) and as allelic forms CD45.1/Ly-5.1 and CD45.2/Ly-5.2. The former

arises as a result of alternative splicing of different exons. The specificity of antibodies to CD45.1 and CD45.2 has been exploited in studies where resolution of donor and recipient cells is essential, such as in gene therapy or bone marrow transplantation. (Ref.3,4)

Synonyms:

PTPRC, Leukocyte common antigen, L-CA, T200