

Product datasheet for AM08081FC-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

MHC Class I H2 Kd/Dd Mouse Monoclonal Antibody [Clone ID: 34-1-2S]

Product data:

Product Type: Primary Antibodies

Clone Name: 34-1-2S

Applications: FC

Recommended Dilution: Flow Cytometry: < / = 1 μ g/10e6 cells.

Reactivity: Mouse
Host: Mouse
Isotype: IgG2a

Clonality: Monoclonal

Immunogen: C3HalphaBDF1 mouse splenocytes.

Specificity: This antibody is specific to an epitope in the alpha 3 domain that is common to H-2Kd and H-

2Dd. It binds to a common determinant in the alpha 3 domains of H-2Kd and H-2Dd in the

presence or absence of beta 2 Microglobulin. (Ref.7,8) It cross reacts with the alpha 3 domain of H-2Kb. (Ref.1,2)

Formulation: PBS containing 0.09% Sodium Azide as preservative.

Label: FITC

State: Liquid purified Ig fraction.

Label: Fluorescein Isothiocyanate Isomer 1

Concentration: lot specific

Conjugation: FITC

Storage: Store the antibody undiluted at 2-8°C for one month or in (aliquots) at -20°C for longer.

This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.







Background:

The 'classical' MHC Class I molecules are histocompatibility antigens encoded by the H-2 gene complex and consist of heterodimers of highly polymorphic alpha chains noncovalently associated with the invariant beta 2-Microglobulin. (Ref.3,4) These antigens are expressed on most nucleated cells but expression varies on different cell types. MHC Class I molecules present endogenously synthesized peptides to CD8+ T lymphocytes, which are usually cytotoxic T cells. (Ref.5) MHC Class I antigens expressed on thymic epithelial cells regulate the positive and negative selection of CD8+ T cells during T cell ontogeny. (Ref.3,6)