

Product datasheet for AM08082BT-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

H2-D1 Mouse Monoclonal Antibody [Clone ID: 27-11-13S]

Product data:

Product Type: Primary Antibodies

Clone Name: 27-11-13S

Applications: FC

Recommended Dilution: Flow Cytometry: < / = 1 μ g/10e6 cells. (Ref.1-3)

Immunohistochemistry (Acetone-Fixed, Frozen Tissue Sections only).

Reactivity:MouseHost:MouseIsotype:IgG2a

Clonality: Monoclonal

Immunogen: C3H alphaBDF1 splenocytes. (Ref.1)

Specificity: This antibody recognizes the alpha 3 domain of H-2Db class I MHC antigen. It cross-reacts

with the alpha 3 domain of H-2Ld, H-2Dq and H-2Lq, but not H-2Kd or H-2Dd.

Reactivity with haplotypes k, f, p, r, and s has not been observed.

Customer feedback: Clone 27-11-13S does not cross-react with human cells.

Formulation: PBS containing 0.09% Sodium Azide as preservative.

Label: Biotin

State: Liquid purified Ig fraction.

Concentration: lot specific
Conjugation: Biotin

Storage: Store the antibody undiluted at 2-8°C for one month or in (aliquots) at -20°C for longer.

This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: histocompatibility 2, D region locus 1

Database Link: Entrez Gene 14964 Mouse

P01899





H2-D1 Mouse Monoclonal Antibody [Clone ID: 27-11-13S] - AM08082BT-N

Background:

In the mouse the MHC Class 1 loci are called H2K, H2L and H2D. The equivalent loci in the

human MHC are HLA (Human Leukocyte Antigen).

MHC class I molecules are heterodimers, consisting of a single transmembrane polypeptide chain (the a-chain) which is highly polymorphic, and the invariant beta 2 Microglobulin (which is encoded elsewhere, not in the MHC). MHC class I molecules are found on almost every nucleated cell of the body. Their major function is to present peptide fragments derived from

antigens to cytotoxic T cells.

Synonyms:

H-2D(B), H2-D1