

Product datasheet for AM08084PU-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Cd86 Rat Monoclonal Antibody [Clone ID: 2D10]

Product data:

Product Type: Primary Antibodies

Clone Name: 2D10

Applications: FC, FN, IHC, IP **Recommended Dilution: ELISA.** (Ref.4)

Flow Cytometry: Identification and enumeration of CD86+ cells. (Ref.1,2,4,6,7)

Immunoprecipitation (Ref.1)

Immunohistochemistry on Frozen Sections: Identification of CD86+ cells. (Ref.4).

In vivo and *in vitro* blocking assays. (Ref.1-3,5)

Reactivity: Mouse

Host: Rat

Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Mouse B cell lymphoma cell line 5C2.

Specificity: This antibody recognises CD86, the B7-2 costimulatory molecule and a ligand for CD28 and

CD152.

Formulation: 100 mM Borate Buffered Saline, pH 8.2.

No preservatives or amine-containing buffer salts added.

State: Purified

State: Liquid purified Ig fraction.

Concentration: lot specific

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: CD86 antigen

Database Link: Entrez Gene 12524 Mouse

P42082





Cd86 Rat Monoclonal Antibody [Clone ID: 2D10] - AM08084PU-N

Background:

CD86, also known as B72, is a type I transmembrane glycoprotein and a member of the immunoglobulin superfamily of cell surface receptors. It is expressed at high levels on resting peripheral monocytes and dendritic cells and at very low density on resting B and T lymphocytes. CD86 expression is rapidly upregulated by B cell specific stimuli with peak expression at 18 to 42 hours after stimulation. CD86, along with CD80/B71, is an important accessory molecule in T cell costimulation via its interaction with CD28 and CD152/CTLA4. Since CD86 has rapid kinetics of induction, it is believed to be the major CD28 ligand expressed early in the immune response. It is also found on malignant Hodgkin and Reed Sternberg (HRS) cells in Hodgkin's disease. CD86 interacts with HHV8 (KSHV) MIR2 protein.

Synonyms:

CD28LG2, Activation B7-2 antigen, B70, BU63, B7.2, FUN-1