

Product datasheet for AM08089FC-N

Icam2 Rat Monoclonal Antibody [Clone ID: 3C4]

Product data:

Product Type: Primary Antibodies

Clone Name: 3C4
Applications: FC

Recommended Dilution: Flow Cytometry: < / = 1 μ g/10e6 cells. (Ref.1,2)

Reactivity: Mouse
Host: Rat
Isotype: IgG2a

Clonality: Monoclonal

Immunogen: COS cells transfected with mouse ICAM-2 cDNA. (Ref.1)

Specificity: This antibody is specific to ICAM-2 (CD102).

It blocks the interaction between ICAM-2 and LFA-1. (Ref.1)

Formulation: PBS containing 0.09% Sodium Azide as preservative.

Label: FITC

State: Liquid purified Ig fraction.

Label: Fluorescein Isothiocyanate Isomer 1

Concentration: lot specific

Conjugation: FITC

Storage: Store the antibody undiluted at 2-8°C for one month or in (aliquots) at -20°C for longer.

This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: intercellular adhesion molecule 2

Database Link: Entrez Gene 15896 Mouse

P35330



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Icam2 Rat Monoclonal Antibody [Clone ID: 3C4] - AM08089FC-N

Background:

ICAM2 is a member of the intercellular adhesion molecule (ICAM) family. All ICAM proteins are type I transmembrane glycoproteins, contain 2-9 immunoglobulin like C2 type domains, and bind to the leukocyte adhesion LFA1 protein. This protein may play a role in lymphocyte recirculation by blocking LFA1 dependent cell adhesion. It mediates adhesive interactions important for antigen specific immune response, NK cell mediated clearance, lymphocyte recirculation, and other cellular interactions important for immune response and surveillance.

Synonyms:

ICAM-2