

Product datasheet for AM08109PU-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD21 (CR2) Mouse Monoclonal Antibody [Clone ID: BB6-11C9.6]

Product data:

Product Type: Primary Antibodies

Clone Name: BB6-11C9.6
Applications: FC, IHC, IP

Recommended Dilution: Flow Cytometry. (Ref.2-3)

Immunohistochemistry (Acetone-Fixed, Frozen Tissue Sections only). (Ref.2)

Immunoprecipitation. (Ref.2)

Reactivity: Porcine
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Specificity: This antibody is specific to Porcine CD21 (Mr. 160 kDa).

It stains the following percentages of surface immunoglobulin (slg) positive cells in porcine lymphoid tissues: 92-96% in peripheral blood; 74-98% in mesenteric lymph nodes; 97-100% in ileal Peyer's patches (ILPP); 93-97% in spleen; and 0% in thymus. The antibody does not stain B cells in cord blood or spleen from pig fetuses (105 days gestation). Does not stain B cells in

cord blood or spleen from pig fetuses (105 days gestation).

Formulation: 100 mM Borate Buffered Saline, pH 8.2.

No preservatives or amine-containing buffer salts added.

State: Purified

State: Liquid purified Ig fraction.

Concentration: lot specific

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C.

DO NOT FREEZE!

Stability: Shelf life: one year from despatch.

Gene Name: complement component 3d receptor 2

Database Link: Entrez Gene 1380 Human

P20023





CD21 (CR2) Mouse Monoclonal Antibody [Clone ID: BB6-11C9.6] - AM08109PU-N

Background: CD21 also known as complement receptor 2 (CR2), C3d receptor or EBV receptor is a 140 kDa

protein. CD21 is a glycosylated type I transmembrane protein consisting of an extracellular face of a series of 15 or 16 CCP domains. CD21 is the receptor for complement components C3d and iC3b as well as the Epstein-Barr virus (EBV) glycoprotein gp350/220. The soluble CD21 (sCD21) was shown to efficiently trigger CD23 signalling pathways in human monocytes. By inducing release of proinflammatory cytokines and upregulating expression of molecules involved in antigen presentation, sCD21 modulates critical monocyte functions that may be

relevant to allergic and inflammatory disorders.

Synonyms: Complement receptor type 2, CR2, C3DR, C3d receptor, EBV Receptor, Dendritic Cell Marker

Protein Families: Druggable Genome, Transmembrane

Protein Pathways: B cell receptor signaling pathway, Complement and coagulation cascades, Hematopoietic cell

lineage