

Product datasheet for AM08116PU-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD8 Mouse Monoclonal Antibody [Clone ID: CT-8]

Product data:

Product Type: Primary Antibodies

Clone Name: CT-8

Applications: FC, FN, IHC, IP

Recommended Dilution: Flow Cytometry: Identification and enumeration of CD8+cells. (Ref.2)

Immunohistochemistry (Acetone-Fixed, Frozen Tissue Sections only): Identification of CD8+

T cells. (Ref.4)

Immunoprecipitation. (Ref.2)

Reactivity: Chicken
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Specificity: This antibody recognizes alpha chain (34-kDa) of Chicken CD8.

Formulation: 100 mM Borate Buffered Saline, pH 8.2.

No preservatives or amine-containing buffer salts added.

State: Purified

State: Liquid purified Ig fraction.

Concentration: lot specific

Conjugation: Unconjugated

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Database Link: Q0R4I4





CD8 Mouse Monoclonal Antibody [Clone ID: CT-8] - AM08116PU-N

Background:

The CD8 antigen is a cell surface glycoprotein found on most cytotoxic T lymphocytes that mediates efficient cell to cell interactions within the immune system. The CD8 antigen, acting as a coreceptor, and the T cell receptor on the T lymphocyte recognize antigen displayed by an antigen presenting cell (APC) in the context of class I MHC molecules. The functional coreceptor is either a homodimer composed of two alpha chains, or a heterodimer composed of one alpha and one beta chain. Both alpha and beta chains share significant homology to immunoglobulin variable light chains. CD8 is expressed on most thymocytes and approximately 1/3 of peripheral blood T cells in humans. CD8 alpha/beta heterodimers are expressed only on TCR alpha/beta T cells, whereas CD8 alpha homodimers can be expressed on alpha/beta and gamma/delta T cells and some NK cells in humans. The pattern of expression in other species has not been well documented.

Synonyms:

CD8 alpha chain, CD8A, MAL