

Product datasheet for AM08129BT-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Chicken Lambda light chain Mouse Monoclonal Antibody [Clone ID: L-1]

Product data:

Product Type: Primary Antibodies

Clone Name: L-1

Applications: ELISA, FC, IF

Recommended Dilution: ELISA.

Flow Cytometry: < / = 1 µg/10e6 cells.

Immunofluorescence.

Reactivity: Chicken
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Specificity: This antibody recognizes Lambda light chains of Chicken Immunoglobulin. (Ref.1)

Formulation: PBS containing 0.09% Sodium Azide as preservative.

Label: Biotin

State: Liquid purified Ig fraction.

Concentration: lot specific
Conjugation: Biotin

Storage: Store the antibody undiluted at 2-8°C for one month or in (aliquots) at -20°C for longer.

This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.





Chicken Lambda light chain Mouse Monoclonal Antibody [Clone ID: L-1] - AM08129BT-N

Background:

All five immunoglobulin classes share the same basic four polypeptide chain structure of two heavy-chains and two light chains. There are five heavy chain types, and two light-chain types (Kappa and Lambda) both having a molecular weight of 22.5kDa. Any heavy-chain type can associate with either light-chain type, but on any immunoglobulin molecule both light-chains are of the same type. Kappa and Lambda consist of a variable region and a constant region and can easily be differentiated by the antigenic properties of the constant region. The ratio of Kappa to Lambda is 70:30, the vast majority of which is bound to heavy-chain in immunoglobulin. In normal individuals low levels of free light-chain arepresent in serum (kappa, 1.6-15.2 mg/L; Lambda, 0.4-4.2mg/L), with the occurrence of multiple myeloma or other B-cell malignancies these levels can be greatly elevated and can be found at high levels in the urine (Bence-Jones proteins).

Synonyms:

Ig lambda chain C regions