

Product datasheet for AM08140RP-N

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

CD44 Mouse Monoclonal Antibody [Clone ID: AV6]

Product data:

Product Type: Primary Antibodies

Clone Name: AV6

Applications: FC

Recommended Dilution: Flow Cytometry: < $/ = 0.2 \mu g/10e6$ cells.

Reactivity: Chicken
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Specificity: This antibody is specific to Chicken CD44 (Mr 90 kDa).

It also reacts with IgM+ Turkey B cells and a subset of Turkey lymphocytes.

Formulation: PBS containing 0.09% Sodium Azide as preservative and a stabilizing agent.

Label: PE

State: Liquid purified Ig fraction.

Label: R-Phycoerythrin

Concentration: lot specific

Conjugation: PE

Storage: Store the antibody undiluted at 2-8°C.

DO NOT FREEZE!

This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Database Link: Q9W6S4

Background: CD44 is a type 1 transmembrane glycoprotein also known as Phagocytic Glycoprotein 1 (pgp

1) and HCAM. CD44 is the receptor for hyaluronate and exists as a large number of different isoforms due to alternative RNA splicing. The major isoform expressed on lymphocytes, myeloid cells, and erythrocytes is a glycosylated type 1 transmembrane protein. Other isoforms contain glycosaminoglycans and are expressed on hematopoietic and non

hematopoietic cells. CD44 is involved in adhesion of leukocytes to endothelial cells, stromal

cells, and the extracellular matrix.






Synonyms:

LHR, MDU2, MDU3, MIC4, CDw44, Epican, ECMR-III, HUTCH-I, Heparan sulfate proteoglycan, Hermes antigen, Hyaluronate receptor, PGP-1

Product images:



Immunofluorescent Staining: Chicken peripheral blood mononuclear cells were double stained with Mouse anti-Chicken CD44-PE and Mouse anti-Chicken Beta-2 Microglobulin. Lymphocytes were gated and analyzed on a FACScan (TM) flow cytometer (BDB, San Jose, CA).