

Product datasheet for AM08165RP-N

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

OriGene Technologies, Inc.

EU: info-de@origene.com CN: techsupport@origene.cn

CD95 (FAS) Mouse Monoclonal Antibody [Clone ID: DX2]

Product data:

Product Type: Primary Antibodies

Clone Name: DX2
Applications: FC

Recommended Dilution: Flow Cytometry: 10 µL/10e6 cells.

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Specificity: This antibody recognizes CD95/Fas/Apo-1.

Crosslinking of CD95 by the Monoclonal antibodies DX2 and DX3 delivers an apoptotic signal to Fas-sensitive cells, indicating that these monoclonal antibodies recognize a functional

epitope of CD95. (Ref.1-5)

Formulation: PBS containing 0.09% Sodium Azide as preservative and a stabilizing agent.

Label: PE

State: Liquid purified Ig fraction.

Label: R-Phycoerythrin

Conjugation: PE

Storage: Store the antibody undiluted at 2-8°C.

DO NOT FREEZE!

This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: Fas cell surface death receptor

Database Link: Entrez Gene 355 Human

P25445





Background:

CD95, also known as FAS or APO1, is a 36 kDa cell surface type I membrane glycoprotein with an apparent molecular weight of 44 kDa on SDS PAGE. CD95 is a member of the TNF receptor family, which includes TNFR1, TNFR2, CD27, CD30 and CD40. Binding of CD95 Ligand to CD95 or crosslinking of CD95 by anti CD95 monoclonal antibodies leads to apoptosis of CD95 expressing cells. CD95 belongs to a subgroup of family members that have a death domain (DD) which contains 70 amino acids near the carboxyl terminal region of the molecule. The binding of adaptor molecules to this DD is responsible for transmitting the death signal for apoptosis. Stimulation of CD95 results in aggregation of its DD, leading to the recruitment of FADD and caspase 8 that together with the receptor form the death inducing signaling complex (DISC). CD95/CD95L is involved in the peripheral deletion of activated mature T cells at the end of the immune response and defects in this pathway predispose to autoimmune disorders. CD95 is also involved in killing of targets such as virus infected cells or cancer cells and killing of inflammatory cells at immune privileged sites.

Synonyms: FASLG receptor, Apo-1 antigen, APT1, FAS1, TNFRSF6

Note: Related Products:

CD95 Antibody -Purified (#AM08165PU-N) CD95 Antibody -APC (#AM08165AC-N) CD95 Antibody -FITC (#AM08165FC-N)

CD95 Antibody -Low Endotoxin (#AM08165LE-N)

CD95 Antibody -Biotin (#AM08165BT-N)