

Product datasheet for AM08170FC-N

LAMP2 Mouse Monoclonal Antibody [Clone ID: H4B4]

Product data:

Product Type: Primary Antibodies

Clone Name: H4B4
Applications: FC

Recommended Dilution: Flow Cytometry: 10 µL/10e6 cells.

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Specificity: This antibody recognizes CD107b/LAMP-2.

Formulation: PBS containing 0.09% Sodium Azide as preservative.

Label: FITC

State: Liquid purified Ig fraction.

Label: Fluorescein Isothiocyanate Isomer 1

Conjugation: FITC

Storage: Store the antibody undiluted at 2-8°C for one month or in (aliquots) at -20°C for longer.

This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: lysosomal associated membrane protein 2

Database Link: Entrez Gene 3920 Human

P13473



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



LAMP2 Mouse Monoclonal Antibody [Clone ID: H4B4] - AM08170FC-N

Background:

CD107b, also known as lysosomal-associated membrane protein 2 (LAMP-2), is a heavily glycosylated, type I transmembrane protein that constitutes the major sialoglycoproteins on lysosomal membranes. It is a ligand for galaptin, an S-type lectin present in extracellular matrix, through its recognition of acetyllactosamine oligosaccharide chains, and is a ligand for E-selectin-mediated cell adhesion.

CD107b is expressed by granulocytes, T cells, macrophages, dendritic cells, activated platelets tonsillar epithelium and some tumor cell lines, including U937 and KG1a. It is also a widely expressed intracellular antigen. LAMP-2 may function in protecting the inner surface of the lysosomal membrane by forming a barrier to lysosomal hydrolases. The upregulation of both CD107a and CD107b on the surface of tumor cell lines has been associated with their enhanced metastatic potential, where they may increase adhesion to extracellular matrix and endothelium. (Ref.1-7)

Synonyms:

LAMP-2, LAMP-2C, LAMPB